

BMI may mediate inverse link between fiber intake, knee OA

December 14 2018



(HealthDay)—The inverse association between fiber intake and the risk

for incident symptomatic knee osteoarthritis may be partially mediated by body mass index, according to a study published in the December issue of *The Journal of Nutrition*.

Zhaoli Dai, Ph.D., from the Boston University School of Medicine, and colleagues used data from the Osteoarthritis Initiative (OAI) and the Framingham Offspring Osteoarthritis Study to assess how BMI and inflammation might impact the observed association between greater fiber intake and the lower risk for symptomatic [knee osteoarthritis](#). Participants were followed for four years and 9.5 years, respectively, in the two studies. Incidence of symptomatic knee osteoarthritis was defined as new onset of a combination of knee pain and radiographic osteoarthritis.

The researchers found that symptomatic knee osteoarthritis developed in 861 knees among 2,876 individuals in the OAI study and in 143 knees among 971 individuals in the Framingham study. In OAI, for participants with higher fiber intake (≥ 21 g/day versus

Citation: BMI may mediate inverse link between fiber intake, knee OA (2018, December 14) retrieved 25 January 2023 from <https://medicalxpress.com/news/2018-12-bmi-inverse-link-fiber-intake.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--