

Opioid use and misuse following treatment for head and neck cancer

26 December 2018, by Garth Sundem



Jessica McDermott, M.D., and colleagues show 7 percent opioid use six months after head and neck cancer treatment ends. Credit: University of Colorado Cancer Center

Cancer patients are often prescribed pain medications, for example during recovery from surgical procedures. However, for many cancer patients, the use of opioid pain medications during treatment can be a gateway to misuse or addiction once treatment ends. Now with cancer patients living longer than ever before, protecting quality of life in the months, years, or decades after treatment is becoming increasingly important, including guarding against the risk of opioid addiction.

"We felt like it was long term problem for some of our head and neck [cancer patients](#), but didn't know how much of problem," says first author Jessica McDermott, MD, investigator at CU Cancer Center and assistant professor at the CU School of Medicine.

To discover the extent of opioid use and abuse in head and [neck cancer](#) patients, McDermott and colleagues searched the SEER/Medicare database to identify 976 patients treated between 2008 and 2011 for oral or oropharynx cancer. In all, 811 of these patients received prescriptions for opioid [pain medications](#) during treatment. Three months after treatment ended, 150 of these patients continued to have active opioid prescriptions. Six months after treatment, 68 patients or 7 percent of the total population continued to use opioid pain medication. Results are published online ahead of print in the journal *Otolaryngology-Head and Neck Surgery*.

"You shouldn't need opioids at the six-month point," McDermott says. "We hope that we can use this data to help patients manage pain better."

In fact, McDermott suggests that because younger patients in this (and other) studies were at greater risk for opioid misuse, and the SEER/Medicare database is heavily skewed to include data from [older patients](#), the true percent of [opioid misuse](#) in the overall head and neck cancer patient population is likely higher than the current study suggests. Additional risk factors for continued opioid use included opioid prescription prior to cancer treatment, and a history of smoking and/or alcohol use.

Interestingly, patients prescribed oxycodone as their first opiate were less likely to continue use at 3 and 6 months after treatment, than patients initially prescribed hydrocodone or other opiates including fentanyl, hydromorphone, meperidine hydrochloride, morphine, nalbuphine, or tramadol.

"We don't know why this is the case, but we think that maybe patients know the word 'oxycodone' and are more aware of the potential for addiction than they might be if prescribed a less well-known drug or one they consider less dangerous like hydrocodone," McDermott says.

The group sees the current study as a way to understand the ecosystem of pain control and [opioid patients](#), with the goal of working to change doctors' strategies for pain control.

"If a patient needed opioids for pain, I wouldn't keep them away, but especially if they have risk factors, I might counsel them more about the risks of addiction and misuse, and keep an eye on it," McDermott says.

More information: Jessica D. McDermott et al, Short- and Long-term Opioid Use in Patients with Oral and Oropharynx Cancer, *Otolaryngology–Head and Neck Surgery* (2018). [DOI: 10.1177/0194599818808513](https://doi.org/10.1177/0194599818808513)

Provided by CU Anschutz Medical Campus

APA citation: Opioid use and misuse following treatment for head and neck cancer (2018, December 26) retrieved 2 May 2021 from <https://medicalxpress.com/news/2018-12-opioid-misuse-treatment-neck-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.