

# Frequent use of aspirin can lead to increased bleeding

22 January 2019



Credit: CC0 Public Domain

A new study published today in the *Journal of the American Medical Association (JAMA)* has found that taking aspirin on a regular basis to prevent heart attacks and strokes, can lead to an increase risk of almost 50% in major bleeding episodes.

The [systematic review](#) from scientists at King's College London and King's College Hospital looked at the overall effects on patients who did not have known cardiovascular disease. They found that while it was associated with a lower risk of heart attacks and other cardiovascular events, it did lead to an increased risk of major bleeding.

While aspirin is known to reduce risks for those who have previously suffered strokes and heart attacks, the evidence of the role of aspirin in the initial prevention of cardiovascular events, is uncertain.

This study looked at the outcomes of trials enrolling more than 1,000 participants with no known history of cardiovascular diseases and which included a follow-up after twelve months. Participants included ones who took aspirin and other who took a placebo or had no treatment at

all.

The results showed that:

- Aspirin use was associated with an 11% lower risk of cardiovascular events.
- Approximately 250 patients needed to be treated with aspirin for 5 years to prevent a single [heart attack](#), stroke or cardiovascular death.
- Aspirin use was associated with a 43% of major bleeding events, compared to those who did not take it.
- Approximately one in 200 people treated with aspirin would have a major bleed.
- No effect was seen with aspirin on new cancer diagnoses or deaths.

Lead author, Dr. Sean Zheng, Academic Clinical Fellow in Cardiology at King's College London said: "This study demonstrates that there is [insufficient evidence](#) to recommend routine [aspirin](#) use in the prevention of heart attacks, strokes and cardiovascular deaths in people without cardiovascular disease.

"There has been more uncertainty surrounding what should be done in patients who are at higher risk of [cardiovascular disease](#) and in patients with diabetes. This study shows that while [cardiovascular events](#) may be reduced in these [patients](#), these benefits are matched by an increased risk of major bleeding events.

"Aspirin use requires discussion between the patient and their physician, with the knowledge that any small potential cardiovascular benefits are weighed up against the real risk of severe bleeding."

Provided by King's College London

APA citation: Frequent use of aspirin can lead to increased bleeding (2019, January 22) retrieved 14 April 2021 from <https://medicalxpress.com/news/2019-01-frequent-aspirin.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*