


Sleep deprivation accelerates Alzheimer's brain damage

24 January 2019



Credit: CC0 Public Domain

Poor sleep has long been linked with Alzheimer's disease, but researchers have understood little about how sleep disruptions drive the disease.

Now, studying [mice](#) and people, researchers at Washington University School of Medicine in St. Louis have found that [sleep deprivation](#) increases levels of the key Alzheimer's [protein tau](#). And, in follow-up studies in the mice, the research team has shown that sleeplessness accelerates the spread through the brain of toxic clumps of tau - a harbinger of brain damage and decisive step along the path to dementia.

These findings, published online Jan. 24 in the journal *Science*, indicate that lack of sleep alone helps drive the disease, and suggests that good sleep habits may help preserve brain health.

"The interesting thing about this study is that it suggests that real-life factors such as sleep might affect how fast the disease spreads through the brain," said senior author David Holtzman, MD, the Andrew B. and Gretchen P. Jones Professor and head of the Department of Neurology. "We've known that sleep problems and Alzheimer's are

associated in part via a different Alzheimer's protein—amyloid beta—but this study shows that sleep disruption causes the damaging protein tau to increase rapidly and to spread over time."

Tau is normally found in the brain—even in [healthy people](#)—but under certain conditions it can clump together into tangles that injure nearby tissue and presage cognitive decline. Recent research at the School of Medicine has shown that tau is high in [older people](#) who sleep poorly. But it wasn't clear whether lack of sleep was directly forcing tau levels upward, or if the two were associated in some other way. To find out, Holtzman and colleagues including first authors Jerrah Holth, Ph.D., a staff scientist, and Sarah Fritschi, Ph.D., a former postdoctoral scholar in Holtzman's lab, measured tau levels in mice and people with normal and disrupted sleep.

Mice are nocturnal creatures. The researchers found that tau levels in the fluid surrounding brain cells were about twice as high at night, when the animals were more awake and active, than during the day, when the mice dozed more frequently. Disturbing the mice's rest during the day caused daytime tau levels to double.

Much the same effect was seen in people. Brendan Lucey, MD, an assistant professor of neurology, obtained cerebrospinal fluid—which bathes the brain and spinal cord—from eight people after a normal night of sleep and again after they were kept awake all night. A sleepless night caused tau levels to rise by about 50 percent, the researchers discovered.

Staying up all night makes people stressed and cranky and likely to sleep in the next chance they get. While it's hard to judge the moods of mice, they, too, rebounded from a sleepless day by sleeping more later. To rule out the possibility that stress or behavioral changes accounted for the changes in tau levels, Fritschi created genetically modified mice that could be kept awake for hours at

a time by injecting them with a harmless compound. **More information:** J.K. Holth et al., "The sleep-wake cycle regulates brain interstitial fluid tau in mice and CSF tau in humans," *Science* (2019). [science.sciencemag.org/lookup/...1126/science.aav2546](https://science.sciencemag.org/lookup/doi/10.1126/science.aav2546)

Using these mice, the researchers found that staying awake for prolonged periods causes tau levels to rise. Altogether, the findings suggest that tau is routinely released during waking hours by the normal business of thinking and doing, and then this release is decreased during sleep allowing tau to be cleared away. Sleep deprivation interrupts this cycle, allowing tau to build up and making it more likely that the protein will start accumulating into harmful tangles.

Provided by Washington University School of Medicine

In people with Alzheimer's disease, tau tangles tend to emerge in parts of the brain important for memory—the hippocampus and entorhinal cortex—and then spread to other brain regions. As tau tangles mushroom and more areas become affected, people increasingly struggle to think clearly.

To study whether the spread of tau tangles is affected by sleep, the researchers seeded the hippocampi of mice with tiny clumps of tau and then kept the animals awake for long periods each day. A separate group of mice also was injected with tau tangles but was allowed to sleep whenever they liked. After four weeks, tau tangles had spread further in the sleep-deprived mice than their rested counterparts. Notably, the new tangles appeared in the same areas of the [brain](#) affected in people with Alzheimer's.

"Getting a good night's sleep is something we should all try to do," Holtzman said. "Our brains need time to recover from the stresses of the day. We don't know yet whether getting adequate sleep as people age will protect against Alzheimer's disease. But it can't hurt, and this and other data suggest that it may even help delay and slow down the disease process if it has begun."

The researchers also found that disrupted sleep increased release of synuclein protein, a hallmark of Parkinson's disease. People with Parkinson's—like those with Alzheimer's—often have sleep problems.

APA citation: Sleep deprivation accelerates Alzheimer's brain damage (2019, January 24) retrieved 31 July 2022 from <https://medicalxpress.com/news/2019-01-deprivation-alzheimer-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.