


Ultrasound predicts fistula formation for kidney dialysis patients

29 January 2019, by Jeff Hansen



Credit: CC0 Public Domain

For more than 20 years, Michael Allon, M.D., and Michelle Robbin, M.D.—along with Carlton Young, M.D., and interventional radiology colleagues in the University of Alabama at Birmingham School of Medicine divisions of Nephrology and Transplantation, and the Department of Radiology—have collaborated to improve the care of end-stage kidney disease patients who are on hemodialysis.

One vexing problem they face is trying to form fistulas in the arms of kidney-failure patients, in preparation for initiating dialysis.

"People with end-stage renal disease do dialysis for four hours, three days a week," said Robbin, a professor of radiology and chief of Ultrasound. "The problem with fistulas is about 50 percent of them fail to mature, so they are not usable for dialysis. Nobody knows the right answer yet as to how to create a fistula that will work every time."

Hemodialysis requires a high blood-flow rate from the patient to the dialysis machine, where the blood is purified and excess fluid removed. Veins are normally too small and thin to support this high

rate, and arteries deep in the arm are not easy to reach. So in the weeks or months before patients need to begin hemodialysis, surgeons attempt to create an arteriovenous fistula—a surgical connection between an artery and a vein—in the patients' arms. Higher blood pressure from the artery connection helps thicken the walls of the vein; in four to six weeks—if the fistula is successful—the vein can tolerate the stress of dialysis.

According to the National Institutes of Health, 468,000 Americans are kidney dialysis patients.

Over the years, Robbin and Allon, a professor of nephrology with a research focus on clinical dialysis, and their surgical and interventional colleagues have contributed substantially to the evaluation and treatment of [dialysis](#) patients. One example is a recent, multicenter NIH-funded clinical research publication using post-operative ultrasound imaging to predict fistula maturation success.

The researchers followed 602 fistula surgery patients in an observational study—using ultrasound imaging at one day, two weeks and six weeks after surgery. The question was whether some or any of 14 different variables, including seven case-mix factors like age, sex, race and body mass index, correlated with later fistula success.

To their surprise, just three of the variables moderately, but significantly, predicted successful arteriovenous fistula maturation—those three were arteriovenous fistula blood flow, fistula diameter and fistula depth. None of the other factors improved prediction.

By providing probabilities of maturation, the ultrasound measurements can "facilitate decisions by the clinician, such as the need for surgical or percutaneous intervention or abandonment of the arteriovenous fistula," Robbin said. "The richness of

the present data will also be highly valuable for practice guideline workgroups to formulate recommendations, taking into account other factors such as invasiveness of the remedial procedures, burden to the patient, costs and differences in process of care in individual centers."

"It would be ideal," Robbin said, "to tell the surgeon and clinical team, based on the six-week ultrasound, 'this fistula has a high chance of maturation without intervention,' or conversely 'this fistula will not mature without intervention.' But we still don't know all the answers, and further research is needed into the problem of [fistula](#) maturation."

The study, "Prediction of [arteriovenous fistula](#) clinical [maturation](#) from postoperative ultrasound measurements: Findings from the Hemodialysis Fistula Maturation Study," was published online ahead of print in the *Journal of the American Society of Nephrology*.

More information: Michelle L. Robbin et al. Prediction of Arteriovenous Fistula Clinical Maturation from Postoperative Ultrasound Measurements: Findings from the Hemodialysis Fistula Maturation Study, *Journal of the American Society of Nephrology* (2018). [DOI: 10.1681/ASN.2017111225](#)

Provided by University of Alabama at Birmingham
APA citation: Ultrasound predicts fistula formation for kidney dialysis patients (2019, January 29) retrieved 28 May 2021 from <https://medicalxpress.com/news/2019-01-ultrasound-fistula-formation-kidney-dialysis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.