

Vitamin D could lower the risk of developing diabetes

30 January 2019



levels.

Study results appear in the article "Higher serum levels of vitamin D are associated with lower blood glucose levels."

"Although a causal relationship has not been proven, low levels of vitamin D may play a significant role in type 2 diabetes mellitus," says Dr. JoAnn Pinkerton, NAMS executive director. "Vitamin D supplementation may help improve blood sugar control, but intervention studies are still needed."

Credit: CC0 Public Domain

Provided by The North American Menopause Society

The benefits of vitamin D in promoting bone health are already well known. A new study out of Brazil suggests that vitamin D also may promote greater insulin sensitivity, thus lowering glucose levels and the risk of developing type 2 diabetes. Results are published online today in *Menopause*, the journal of The North American Menopause Society (NAMS).

Other recent studies have shown a clear relationship between vitamin D and glycemic control, suggesting that vitamin D increases insulin sensitivity and improves pancreatic beta-cell function. In this cross-sectional study involving 680 Brazilian women aged 35 to 74 years, the goal was to evaluate the possible association between vitamin D deficiency and increased glycemia.

Of the women interviewed, 24 (3.5%) reported using vitamin D supplements. Vitamin D supplementation was found to be negatively associated with high glucose levels. Habitual exposure to the sun also provided the same association, demonstrating that vitamin D deficiencies are associated with high blood glucose



APA citation: Vitamin D could lower the risk of developing diabetes (2019, January 30) retrieved 17 June 2021 from https://medicalxpress.com/news/2019-01-vitamin-d-diabetes.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.