

Vaccine-preventable infections common after peds transplant

11 February 2019



fatality rate was 1.7 percent. Increased risk for hospitalization from a vaccine-preventable [infection](#) was positively associated with age

(HealthDay)—Hospitalization for vaccine-preventable infections occurs in more than 15 percent of pediatric solid organ transplant recipients in the first five years after surgery, according to a study recently published in *JAMA Pediatrics*.

Amy G. Feldman, M.D., from Children's Hospital Colorado in Denver, and [colleagues](#) used data from the Pediatric Health Information System to assess hospitalizations for vaccine-preventable infections among 6,980 patients (mean age at [transplant](#), 8 years) in the first five years after pediatric solid organ transplant.

The researchers found a total of 1,490 cases of vaccine-preventable infections among 1,092 patients (15.6 percent). Of these cases, 13.1 percent occurred during transplant hospitalization. Transplant hospitalizations complicated by vaccine-preventable infections were \$120,498 more expensive than those not complicated by vaccine-preventable infections. For all infections, the case

APA citation: Vaccine-preventable infections common after peds transplant (2019, February 11)
retrieved 11 October 2022 from <https://medicalxpress.com/news/2019-02-vaccine-preventable-infections-common-peds-transplant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.