

Early use of antibiotics in elderly patients with UTIs associated with reduced risk of sepsis

February 27 2019

Prescribing antibiotics immediately for elderly patients with urinary tract infections is linked with a reduced risk of sepsis and death, compared with patients who receive antibiotics in the days following diagnosis, or none at all.

These are the latest findings from researchers at Imperial College London and Public Health England, published in the *BMJ*.

The research team say the results provide further evidence to help GPs make clinical decisions about when to prescribe antibiotics immediately for a [urinary tract infections](#) (UTI) and when to defer treatment to see if symptoms improve on their own, to avoid overuse of antibiotics.

In the research, funded by the National Institute for Health Research, the team looked at records from 157,264 [patients](#) over the age of 65 across England who had been diagnosed by their GP with a suspected or confirmed UTI. Patients had been prescribed antibiotics immediately (87 per cent of cases studied in the research), had antibiotics delayed by up to 7 days (6 per cent of cases), or received no antibiotics at all (7 per cent of cases).

Of the patients who received antibiotics immediately, 0.2 per cent developed sepsis within the following 60 days. After taking into account available information about differences in age, gender, pre-existing

illness and other personal characteristics, the results revealed that compared with patients who received antibiotics immediately, patients who had their antibiotic prescription delayed or received no antibiotics at all were up to eight times more likely to develop sepsis.

The research also revealed that 1.6 per cent of patients who received antibiotics immediately died in the following 60 days. The risk of death over the same time period among patients who had their antibiotic prescription delayed showed a slight increase (16 per cent), while patients who received no antibiotics had over double the risk.

The researchers estimated that, on average, for every 37 patients exposed to no antibiotics and for every 51 patients exposed to deferred antibiotics, one case of sepsis would occur that would not have been seen with immediate antibiotics.

They also found that the rate of hospital admissions roughly doubled (27%) in patients with either no or deferred antibiotic prescriptions, compared with those receiving immediate prescriptions (15%).

Older men, especially those aged over 85 years, and those living in more deprived areas were found to be most at risk.

The researchers stress this study only shows delayed antibiotics are associated with an increased risk of sepsis and death, rather than causing it directly. They add that patients may also have had other [health conditions](#) that the researchers weren't able to account for, which may have contributed to their increased risk of sepsis or death.

Lead author Dr. Myriam Gharbi, from Imperial's School of Public Health, said: "Current national guidelines for GPs recommend they should ask patients about the severity of their symptoms, discuss possible self-care, such as drinking plenty of water to avoid dehydration and

taking paracetamol or ibuprofen for pain relief, and consider a back-up antibiotic prescription to be used if symptoms worsen or have not improved after 48 hours. This is to avoid antibiotic overuse, as sometimes UTIs can get better without medication. However, our research suggests antibiotics should not be delayed in [elderly patients](#)."

UTIs are common in the elderly and may trigger symptoms such as pain when urinating, or needing to use the loo more often. UTIs are most commonly caused by E. coli bacteria, and if not treated the bacteria can trigger blood poisoning.

However, doctors are increasingly concerned about the rise in antibiotic resistance—caused by antibiotics being overprescribed. UTIs are the second most common diagnosis for which antibiotics are prescribed in the UK.

Therefore to help clarify when antibiotics should be prescribed to the elderly with UTIs, the research team studied data from 157, 264 patients aged 65 or above diagnosed with a UTI or suspected UTI, between 2007 and 2015. The data was from the Clinical Practice Research Datalink, which uses anonymised patient data from both GP practices linked to hospital data, allowing the same patients to be tracked between the two settings. The average age of the patients in the study was 77 years old.

Professor Paul Aylin, senior author of the research from the NIHR Health Protection Unit at Imperial, said: "Although antibiotic prescribing must be controlled to help combat the increasing problem of [antibiotic resistance](#), our study suggests early use of antibiotics in elderly patients with UTIs is the safest approach."

Professor Alan Johnson from Public Health England who collaborated on the research said: "Antibiotic resistance is a major threat to public health that is being driven by the [overuse of antibiotics](#). Current

recommendations suggest healthcare professionals take a number of different factors into account when deciding whether to prescribe antibiotics immediately or consider deferring [antibiotics](#) for patients with a suspected urinary tract infection. This study highlights the importance of taking age into account when making clinical decisions about [antibiotic prescribing](#) in order to reduce the risk of complications. This work will help doctors target antibiotic use more effectively and improve patient wellbeing."

More information: "Antibiotic management of urinary tract infection in the elderly in primary care and its association with bloodstream infections and all-cause mortality: a population-based cohort study" *BMJ* (2019). [DOI: 10.1136/bmj.1525](https://doi.org/10.1136/bmj.1525)

Provided by Imperial College London

Citation: Early use of antibiotics in elderly patients with UTIs associated with reduced risk of sepsis (2019, February 27) retrieved 1 February 2024 from <https://medicalxpress.com/news/2019-02-early-antibiotics-elderly-patients-utis.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--