


Behavioral intervention ups physical activity in T2DM

6 March 2019



10.5 metabolic equivalent-hours per week of physical activity volume, 18.9 and 12.5 minutes/day of moderate-to-vigorous intensity physical activity, 4.6 and 3.8 hours/day of light-intensity physical activity, and 10.9 and 11.7 hours/day of [sedentary time](#). Throughout the study, the significant between-group differences were maintained; during the third year, the between-group difference in moderate-to-vigorous intensity physical activity decreased.

"This behavioral intervention strategy was successful in increasing physical activity volume by reallocating sedentary time to light-intensity physical activity and, to a lesser extent, moderate-to-vigorous-intensity [physical activity](#)," the authors write.

Several authors disclosed financial ties to the pharmaceutical industry.

More information: [Abstract/Full Text](#) ([subscription or payment may be required](#))

Copyright © 2019 [HealthDay](#). All rights reserved.

(HealthDay)—A behavioral intervention results in a sustained increase in physical activity and decrease in sedentary time among patients with type 2 diabetes, according to a study published in the March 5 issue of the *Journal of the American Medical Association*.

Stefano Balducci, M.D., from "La Sapienza" University in Rome, and colleagues enrolled 300 physically inactive and sedentary patients with type 2 diabetes to receive a [behavioral intervention](#) or standard of care for three years. One individual theoretical counseling session and eight biweekly theoretical and practical counseling sessions were provided each year to participants in the behavioral [intervention](#) group. In the standard care group, participants received only general physician recommendations.

Participants were followed for a median of 3.0 years. The researchers found that in the behavioral intervention and standard care groups, respectively, participants accumulated 13.8 and

APA citation: Behavioral intervention ups physical activity in T2DM (2019, March 6) retrieved 1 May 2021 from <https://medicalxpress.com/news/2019-03-behavioral-intervention-ups-physical-t2dm.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.