

# Research examines complications during birth and link to later social anxiety in children

20 March 2019



Dr. Santiago Morales, of the University of Maryland.

**More information:** Gabriela L. Suarez et al, Perinatal complications are associated with social anxiety: Indirect effects through temperament, *Infant and Child Development* (2019). DOI: [10.1002/icd.2130](https://doi.org/10.1002/icd.2130)

Provided by Wiley

Credit: CC0 Public Domain

A new study published in *Infant and Child Development* indicates that complications during birth may increase the risk that children will develop social anxiety by their pre-teen years.

For the study, 149 children aged nine to 12 years were screened for behavioral inhibition—a tendency to exhibit a fearful disposition and withdrawal in unfamiliar contexts and situations—and assessed for social anxiety symptoms using parent- and child-reports. Investigators found that perinatal complications were associated with higher levels of behavioral inhibition and social anxiety symptoms.

Additionally, analyses suggested that behavioral inhibition acted as a pathway between birth complications and social anxiety symptoms.

"This study sets the stage for future longitudinal work examining whether childhood temperament is a developmental path by which birth complications lead to [social anxiety](#) symptoms," said lead author

APA citation: Research examines complications during birth and link to later social anxiety in children (2019, March 20) retrieved 30 September 2022 from <https://medicalxpress.com/news/2019-03-complications-birth-link-social-anxiety.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*