

Brain region that produces emotions is larger in vets, service members with mild TBI and PTSD

29 April 2019, by Mike Richman



Dr. Mingxiong Huang is a neuroimaging scientist at the VA San Diego Health Care System. Credit: Kevin Walsh

A new study finds that veterans and active-duty service members with combat-related PTSD and mild traumatic brain injury had larger amygdalas—the region of the brain that processes such emotions as fear, anxiety, and aggression—than those with only brain injuries.

The findings appeared online April 25 in the *Journal of Head Trauma Rehabilitation*.

Through magnetic resonance imaging, the researchers found that the right and left sides of the amygdala in people with combat-related PTSD and [mild traumatic brain injury](#) (mTBI) were larger than those in people with only combat-related mTBI. The amygdala is an almond-shaped section of tissue in the temporal portion of the [brain](#) and is key to triggering PTSD symptoms.

The researchers caution that the findings were based on an observational study and therefore

can't prove a cause-and-effect relationship—only a correlation.

The study included 89 veterans and active-duty military personnel, about a third of whom had both PTSD and mTBI. The rest formed the mild-TBI-only control group. A mild traumatic brain injury is also known as a concussion.

"This is an intriguing structural finding, given the role of the amygdala in the challenging [neuropsychological] symptoms witnessed in casualties of combat-related mTBI and PTSD," the researchers write. "Further investigation is needed to determine whether amygdala size could be used to screen people at risk for PTSD, or whether it could be used to monitor the [effectiveness of medical solutions]."

The study's lead author, Dr. Mingxiong Huang, is a neuroimaging scientist at the VA San Diego Health Care System. He says the finding of a larger amygdala in veterans with combat-related PTSD and mTBI was a bit of a surprise.

"Some previous [PTSD research](#) showed declines in amygdala volume based on the assumption of a loss of size due to injuries," says Huang, also a professor in the department of radiology at the University of California San Diego (UCSD). "Our finding of increased amygdala volume seems to point to different mechanisms, such as an exposure to repetitive fear and stress."

Such exposure, he adds, may lead to an abnormal growth of the neural networks within the amygdala, a development that has been reported in animal studies but hasn't been fully explored in human PTSD studies. More studies involving people with non-combat PTSD are needed to generalize this finding to other types of PTSD, he notes.

A co-author of the paper, Dr. Douglas Chang, is a physician and researcher at VA San Diego.

"The amygdala is involved with processing threat perception and arousal and with linking emotion to experience in complex ways," says Chang, who is also a professor of orthopedic surgery at UCSD. "A larger amygdala volume may be a sign of hyperactivity with an enlarged neural network. But we don't know whether this is an attempt by the brain to cope with PTSD or whether the growth and enlargement is causing symptoms, like an electrical storm."

He adds: "The situation may also resemble scar tissue formation on skin. Is this an organized response by the body to heal itself, or is the scar tissue going haywire and forming a grossly disfigured area? Another possibility is that this study simply identified at-risk people for PTSD with a pre-existing condition: an enlarged amygdala."

Combat-related PTSD and mTBI are leading health care concerns in veterans and service members. It's not unusual for both conditions to occur in the same person, based on evidence from a cross section of studies. Some of the symptoms are similar, such as depression, anxiety, insomnia, fatigue, and changes in memory and concentration. However, the effects of PTSD and mTBI on neural pathways in the brain, as well as the impact of the co-existence of the two, are not fully understood.

Scientists in Huang's study conducted the testing at VA San Diego and at two Marine Corps bases in California. They measured intracranial volume, a key statistic used to analyze the size of the brain and brain regions, especially in cases of neurodegenerative diseases.

The size of the right amygdala was 0.122 percent of total brain volume, on average, in the group with mTBI and PTSD. It was 0.115 percent in the cohort with only mTBI. The size of the left amygdala was 0.118 percent of brain volume in those with mTBI and PTSD, compared with 0.112 percent in the mTBI group. The researchers found both of those differences to be "statistically significant."

The study team also examined the caudate, the

hippocampus, the anterior cingulate cortex, and the cerebral cortex. Those brain regions, like the amygdala, are in the limbic system, which controls basic emotions, including fear, pleasure, and anger. The researchers found no "statistically significant" differences in those regions, suggesting that only changes in the amygdala are linked to PTSD symptoms in people with mTBI, according to Chang.

The study authors say the findings have several implications for research and treatment.

"To be able to see a structural difference between these two cohorts and in this stage of PTSD really points to something going on with the [amygdala](#)," Chang says. "Can we use this as a screening tool to identify people at risk? Maybe this is an adaptive response that we can monitor and use to track different kinds of mental health treatment approaches. Maybe yoga is helpful, maybe mindfulness meditation is helpful, maybe exercise is helpful. Perhaps there are drugs that can protect somebody against these traumas or to help improve their conditions. To be able to identify something that's changed in a quantitative way is amazing. It opens the door to many possibilities to help treat this problem."

More information: Joel Pieper et al, Brain Amygdala Volume Increases in Veterans and Active-Duty Military Personnel With Combat-Related Posttraumatic Stress Disorder and Mild Traumatic Brain Injury, *Journal of Head Trauma Rehabilitation* (2019). [DOI: 10.1097/HTR.0000000000000492](https://doi.org/10.1097/HTR.0000000000000492)

Provided by Veterans Affairs Research Communications

APA citation: Brain region that produces emotions is larger in vets, service members with mild TBI and PTSD (2019, April 29) retrieved 10 November 2022 from <https://medicalxpress.com/news/2019-04-brain-region-emotions-larger-vets.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.