


How electrical stimulation reorganizes the brain

10 June 2019



will translate to non-invasive techniques, such as transcranial magnetic stimulation, and to other patient populations remains to be determined.

More information: Intracortical Dynamics Underlying Repetitive Stimulation Predicts Changes in Network Connectivity, *JNeurosci* (2019). [DOI: 10.1523/JNEUROSCI.0535-19.2019](https://doi.org/10.1523/JNEUROSCI.0535-19.2019)

Provided by Society for Neuroscience

Location of electrodes exhibiting significant stimulation responses and modulation of the stimulation response. Credit: Huang *et al.*, *JNeurosci* (2019)

Recordings of neural activity during therapeutic stimulation can be used to predict subsequent changes in brain connectivity, according to a study of epilepsy patients published in *JNeurosci*. This approach could inform efforts to improve brain stimulation treatments for depression and other psychiatric disorders.

Corey Keller and colleagues delivered [electrical stimulation](#) from implanted electrodes in 14 patients while recording participants' [brain activity](#).

Repeated sets of stimulation resulted in progressive changes to the brain's response to simulation, with stronger responses in brain regions connected to the stimulation site.

The researchers observed these changes in a matter of minutes, suggesting that electrical stimulation induces the brain to rapidly reorganize itself.

Assessing brain activity before, during, and after simulation has the potential to personalize neuromodulation therapies. Whether these results

APA citation: How electrical stimulation reorganizes the brain (2019, June 10) retrieved 17 August 2022 from <https://medicalxpress.com/news/2019-06-electrical-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.