

Late-life diabetes status tied to new cognitive impairment

19 June 2019



shaped association was noted between hemoglobin A1c (HbA1c) and incident dementia. Independent of HbA1c, glycated albumin and fructosamine were also associated with incident dementia. Incident mild [cognitive impairment](#) was associated with HbA1c and fructosamine.

"In [older adults](#) with diabetes, maintaining glycemic control is an important avenue for mitigating cognitive impairments into older age," the authors write.

More information: [Abstract/Full Text](#) ([subscription or payment may be required](#))

Copyright © 2019 [HealthDay](#). All rights reserved.

(HealthDay)—Having diabetes, poor glycemic control, and longer diabetes duration are associated with worse cognitive outcomes in older adults during a median follow-up of five years, according to a study recently published in *Diabetes Care*.

Andreea M. Rawlings, Ph.D., from the Johns Hopkins Bloomberg School of Public Health in Baltimore, and colleagues analyzed data from 5,099 participants (59 percent female; mean age at baseline, 76 years) of the Atherosclerosis Risk in Communities Study. Participants were followed from 2011-2013 until 2016-2017.

The researchers found associations between [diabetes](#) (hazard ratio [HR], 1.14; 95 percent confidence interval [CI], 1.00 to 1.31), [poor glycemic control](#) in persons with diabetes (HR, 1.31; 95 percent CI, 1.05 to 1.63), and longer diabetes duration (at least five years versus less than five years, HR, 1.59; 95 percent CI, 1.23 to 2.07) and incident cognitive impairment. A J-

APA citation: Late-life diabetes status tied to new cognitive impairment (2019, June 19) retrieved 27 April 2021 from <https://medicalxpress.com/news/2019-06-late-life-diabetes-status-tied-cognitive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.