

# Phenols in cocoa bean shells may reverse obesity-related problems in mouse cells

June 21 2019, by Sharita Forrest



Compounds in cocoa bean shells fight obesity, inflammation and insulin resistance in mouse cells, according to new research by University of Illinois food science professor Elvira Gonzalez de Mejia and Miguel Rebello-Hernanz, a visiting scholar in food science. Credit: Fred Zwicky

Scientists may have discovered more reasons to love chocolate.

A new study by researchers at the University of Illinois suggests that three of the phenolic compounds in [cocoa](#) bean shells have powerful effects on the fat and [immune cells](#) in mice, potentially reversing the [chronic inflammation](#) and insulin resistance associated with obesity.

Visiting scholar in food science Miguel Rebollo-Hernanz and Elvira Gonzalez de Mejia, a professor in the department, found that cocoa shells contain high levels of three beneficial bioactive chemicals also found in cocoa, coffee and green tea—protocatechuic acid, epicatechin and procyanidin B2.

Rebollo-Hernanz, the study's lead author, created a water-based extract containing these compounds and tested its effects on white fat cells called adipocytes and immune cells called macrophages. Using computer modeling and bioinformatic techniques, he also examined the impact that each of the phenolics individually had on the cells.

"The objectives of the study were to test whether the bioactive compounds in the cocoa shells were efficacious against macrophages—the [inflammatory cells](#)—at eliminating or reducing the biomarkers of inflammation," said de Mejia, also a director of nutritional sciences. "We wanted to see if the phenolics in the extract blocked or reduced the damage to fat cells' [mitochondria](#) and prevented insulin resistance."

Similar to batteries within cells that burn fat and glucose to generate energy, mitochondria can become damaged when high levels of fat, glucose and inflammation occur in the body, de Mejia said.

When the scientists treated adipocytes with the aqueous extract or the three phenolic compounds individually, damaged mitochondria in the cells were repaired and less fat accumulated in the adipocytes, blocking inflammation and restoring the cells' insulin sensitivity, Rebollo-Hernanz



said.

The scientists reported their findings recently in a paper published in the journal *Molecular Nutrition and Food Research*.

When adipocytes accumulate too much fat, they promote the growth of macrophages. This initiates a toxic cycle in which the adipocytes and macrophages interact, emitting toxins that inflame fat tissue, de Meija said.



Cocoa shells, a waste byproduct of roasting cocoa beans to produce chocolate, contain significant amounts of three healthful bioactive compounds that are also found in cocoa, coffee and tea. Credit: Fred Zwicky

Over time, this chronic inflammation impairs cells' ability to take up glucose, leading to [insulin resistance](#) and possibly type 2 diabetes as glucose levels in the blood escalate.

To recreate the inflammatory process that occurs in the body when macrophages and adipocytes begin their toxic dance, Rebollo-Hernanz grew adipocytes in a solution in which macrophages had been cultured.

"That's when we observed that these inflammatory conditions in the solution increased the [oxidative damage](#)" to the fat cells' mitochondria, he said.

Fewer mitochondria were present in the adipocytes that were grown in the solution, and the mitochondria that did exist in these [cells](#) were damaged, he found.

When the scientists treated the adipocytes with the phenolics in the extract, however, the adipocytes underwent a process called browning, in which they differentiated—that is, converted—from white adipocytes into another form called beige adipocytes.

Beige adipocytes are a specialized form of fat tissue with greater numbers of mitochondria and enhanced fat-burning efficiency.

"We observed that the extract was able to maintain the mitochondria and their function, modulating the inflammatory process and maintaining the adipocytes' sensitivity to insulin," Rebollo-Hernanz said. "Assuming that these phenolics were the main actors in this extract, we can say that consuming them could prevent mitochondrial dysfunction in adipose tissue."

Cocoa shells are a waste byproduct that's generated when cocoa beans are roasted during chocolate production. About 700,000 tons of the

shells are discarded annually, causing environmental contamination if not disposed of responsibly, de Mejia said.

In addition to providing cocoa producers with another potential revenue stream, processing the shells to extract the nutrients would reduce the environmental toxicants generated currently by cocoa [shell](#) waste, de Mejia said.

Once extracted from cocoa bean shells, the [phenolic compounds](#) could be added to foods or beverages to boost products' nutritional value, she said.

**More information:** Miguel Rebollo-Hernanz et al, Cocoa Shell Aqueous Phenolic Extract Preserves Mitochondrial Function and Insulin Sensitivity by Attenuating Inflammation between Macrophages and Adipocytes In Vitro, *Molecular Nutrition & Food Research* (2019). [DOI: 10.1002/mnfr.201801413](#)

Provided by University of Illinois at Urbana-Champaign

Citation: Phenols in cocoa bean shells may reverse obesity-related problems in mouse cells (2019, June 21) retrieved 23 November 2023 from <https://medicalxpress.com/news/2019-06-phenols-cocoa-bean-shells-reverse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.