


Vaccination programs substantially reduce HPV infections and precancerous cervical lesions

27 June 2019



Electron micrograph of a negatively stained human papilloma virus (HPV) which occurs in human warts. Credit: public domain

Human papilloma virus (HPV) vaccination programs have substantially reduced the number of infections and precancerous cervical lesions caused by the virus, according to a study published today in *The Lancet* by researchers from Université Laval and the CHU de Québec-Université Laval Research Centre. Results are so promising that it has now become possible to envision eliminating cervical cancer as a public health problem in the coming decades.

Infections caused by HPV are among the most common sexually transmitted diseases. Certain forms of the virus can lead to anogenital warts while others cause lesions that can develop into [cancer](#) of the mouth, throat, vagina, vulva, anus, or penis, and particularly the cervix. "HPV is found in almost 100% of [cervical cancer](#) cases," states lead author Mélanie Drolet.

The research team headed by Université Laval's Faculty of Medicine Professor Marc Brisson

performed a [meta-analysis](#) of 65 studies in 14 [different countries](#) that have set up HPV vaccine programs in the last ten years. With data on 60 million people, the researchers compared the frequency of HPV infections, anogenital warts, and [precancerous cervical lesions](#) before and after the programs were launched.

Their analysis shows that infections dropped by 83% among girls aged 13 to 19 and 66% among women aged 20 to 24. For anogenital warts, the drop was 67% among 15- to 19-year-old girls, 54% for women aged 20 to 24, and 31% for those aged 25 to 29. Precancerous cervical lesions also dropped by 51% among teens aged 15 to 19 and 31% among women aged 20 to 24.

Vaccination of young women is also producing herd protection for young men. Anogenital warts among males have dropped 48% for those 15 to 19 and 32% for those aged 20 to 24.

Cervical cancer currently affects 1 woman in 150 in Canada, and the 5-year mortality rate is about 25%. "HPV vaccination is still too recent to directly measure its effects on cervical cancer as it can take decades to develop," explains Mélanie Drolet. "However, our analyses show that vaccination is producing substantial reductions in the infections that cause cervical cancer and precancerous lesions. These reductions are a first sign that vaccination could eventually lead to the elimination of cervical cancer as a public health problem. We are now trying to determine when elimination could be achieved and which vaccination and screening programs could help us achieve it faster."

More information: Mélanie Drolet et al, Population-level impact and herd effects following the introduction of human papillomavirus vaccination programmes: updated systematic

review and meta-analysis, *The Lancet* (2019). DOI:
[10.1016/S0140-6736\(19\)30298-3](https://doi.org/10.1016/S0140-6736(19)30298-3)

Provided by Laval University

APA citation: Vaccination programs substantially reduce HPV infections and precancerous cervical lesions (2019, June 27) retrieved 23 April 2021 from

<https://medicalxpress.com/news/2019-06-vaccination-substantially-hpv-infections-precancerous.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.