

Diet quality may affect risk of frailty in older adults

3 July 2019



Prospective Associations of Diet Quality With Incident Frailty in Older Adults: The Health, Aging, and Body Composition Study, *Journal of the American Geriatrics Society* (2019). DOI: 10.1111/jgs.16011

Provided by Wiley

Credit: CC0 Public Domain

Poorer overall diet quality was linked with an increased risk of becoming frail in a study of U.S. community-dwelling older adults, published in the *Journal of the American Geriatric Society*. The quality of the overall diet appeared to be more important than protein intake for a lower risk of frailty.

During 4 years of follow-up, 277 of the 2154 participants—who were 70-81 years and characterized as "robust" or "pre-frail" at the start of the study—developed frailty. Poor- and mediumquality diets were associated with a 92% and 40% higher incidence of frailty compared with goodquality diets, respectively. No association for protein intake was observed.

"The role of single nutrients such as protein in the development of frailty is not fully understood and definitely needs further investigation. Nevertheless, this study contributes to the prevailing idea that the overall quality is important anyway," said lead author Linda Hengeveld, of Vrije Universiteit Amsterdam, in the Netherlands.

More information: Linda M. Hengeveld et al,



APA citation: Diet quality may affect risk of frailty in older adults (2019, July 3) retrieved 30 May 2021 from <u>https://medicalxpress.com/news/2019-07-diet-quality-affect-frailty-older.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.