

Does use of headgear reduce the rate injuries in high school women's lacrosse?

14 July 2019

Headgear worn during women's lacrosse practice and games can reduce the rate of head and face injuries as well as concussions, according to research presented by researchers in the Department of Orthopedics at the New York University Langone Health. The research was presented today at the American Orthopedics Society of Sports Medicine Annual Meeting.

"There has been continued controversy regarding whether or not [headgear](#) use in women's lacrosse will increase or decrease the rate of head injuries," said Dr. Samuel L. Baron, the lead researcher. In 2017, the Public Schools Athletic of New York City became the first high school organization in the country to mandate ASTM standard F3137 headgear for all women's lacrosse players.

The researchers set up a prospective cohort that included eight varsity and junior varsity women's lacrosse teams, as well as their game opponents, who were mandated to wear F3137 headgear for all practice and game events over the course of the 2017 and 2018 seasons. Certified athletic trainers assessed and documented all injuries that occurred as a result of participation on the lacrosse teams and athlete exposures were estimated based on the number of team practice and game events. Injury rates were compared with those from the High School RIO (Reporting Information Online) injury data reports from the 2009 to 2016 seasons.

Over the study period, 17 total injuries were reported during 22,397 exposures for an injury rate of 0.76 injuries per 1,000 athlete-exposures. Two head/face injuries, both of which were classified as concussions, were reported during the study for a head/face injury rate and [concussion](#) rate of 0.09 per 1,000 athlete-exposures. The headgear cohort demonstrated significant decreases in rates of head/face injury in-game concussion and practice non-head/face injury when compared to the control cohort.

Baron and his team concluded that mandated use of F3137 headgear was shown to be effective at lowering the rate of head or face injury and concussions in women's lacrosse. Additionally, mandated headgear use was also shown to lower the rate of [injury](#) to body locations other than the head or face during practice.

Provided by American Orthopaedic Society for Sports Medicine

APA citation: Does use of headgear reduce the rate injuries in high school women's lacrosse? (2019, July 14) retrieved 15 August 2022 from <https://medicalxpress.com/news/2019-07-headgear-injuries-high-school-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.