

# New study finds both components of blood pressure predict heart attack, stroke risk

17 July 2019



Credit: CC0 Public Domain

Both numbers in a blood pressure reading—the "upper" systolic and the "lower" diastolic—Independently predicted the risk of heart attack or stroke in a very large Kaiser Permanente study that included more than 36 million blood pressure readings from more than 1 million people. The study, which was published today in the *New England Journal of Medicine*, runs counter to decades of previous research indicating that high systolic blood pressure is more likely than diastolic pressure to result in adverse outcomes.

"This research brings a large amount of data to bear on a basic question, and it gives such a clear answer," said lead author Alexander C. Flint, MD, Kaiser Permanente stroke specialist and adjunct researcher with the Division of Research. "Every way you slice the data, the systolic and diastolic pressures are both important."

The current retrospective study is "the largest by far of its kind," Dr. Flint said, reviewing 36 million [blood pressure readings](#) taken during outpatient visits between 2007 and 2016, from 1.3 million adult Kaiser Permanente members in Northern

California.

Systolic pressure—the upper number in a blood pressure reading—measures how hard the heart pumps blood into arteries. Diastolic pressure—the bottom number—indicates the pressure on the arteries when the heart rests between beats.

Decades of research have shown that high systolic blood pressure is more likely to result in adverse outcomes. As a result, cardiology guidelines and risk estimation tools focus on the upper number, with some experts arguing that the diastolic number might reasonably be ignored, Flint said.

After adjusting the data for possible confounding factors, the researchers found that while systolic pressure has a greater impact, both systolic and diastolic pressures strongly influenced the risk of heart attack or stroke, regardless of the definition used for [high blood pressure](#) (140/90 mm Hg versus 130/80 mm Hg).

Dr. Flint said the finding that systolic and diastolic hypertension have similar impacts on risk at the lower threshold of 130/80 provides independent support for recent changes that were made in the American College of Cardiology and American Heart Association guidelines, which recommended tighter blood pressure control in higher risk patients with hypertension. The results are also in keeping with findings from the National Institutes of Health's Systolic Blood Pressure Intervention Trial, known as SPRINT.

"Controversy has long persisted about whether [systolic blood pressure](#), diastolic blood pressure, or both contribute to cardiovascular risk," said senior author Deepak L. Bhatt, MD, MPH, executive director of Interventional Cardiovascular Services at Brigham and Women's Hospital and professor of medicine at Harvard Medical School. "This analysis using a very large amount of longitudinal data convincingly demonstrates that both are important,

and it shows that in people who are otherwise generally healthy, lower [blood](#) pressure numbers are better."

Provided by Kaiser Permanente

APA citation: New study finds both components of blood pressure predict heart attack, stroke risk (2019, July 17) retrieved 18 April 2021 from <https://medicalxpress.com/news/2019-07-components-blood-pressure-heart.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*