


Individuals with obesity get more satisfaction from their food

30 July 2019



Change in taste perceptions of chocolate by body mass index (BMI) category. As the number of chocolate samples consumed increased, participants reported lower taste perceptions, on average. Obese individuals tended to report higher taste perceptions compared with normal-weight or overweight individuals ($P=0.02$). Credit: *Journal of the Academy of Nutrition and Dietetics*

The propensity to overeat may, in part, be a function of the satisfaction derived from eating. A new study in the *Journal of the Academy of Nutrition and Dietetics*, published by Elsevier, found no significant difference in taste perceptions between participants of normal weight and those who were overweight. However, participants with obesity had initial taste perceptions that were greater than participants who were not obese, which declined at a more gradual rate than participants who were not obese. This quantification of satisfaction from food may help explain why some people eat more than others.

"Obesity is a major public-health problem. Thirty percent of the US population is obese, and obesity-related health problems (diabetes, hypertension,

etc.) are increasing. Causes of obesity are varied, but food consumption decisions play an important role, especially decisions about what foods to eat and how much to consume. Taste perceptions may lead to overeating. If people with obesity have different taste perceptions than nonobese people, it could lead to better understanding of obesity and possibly designing new approaches to prevent obesity," explained lead investigator Linnea A. Polgreen, Ph.D., Department of Pharmacy Practice and Science, University of Iowa, Iowa City, IA, USA.

As individuals consume more of a food item, they experience diminishing marginal taste perception, which means their level of perceived taste from additional consumption may tend to decline (ie, additional consumption may become less pleasurable). The relationship between perceived taste and quantity consumed has traditionally been referred to as sensory-specific satiety.

In order to determine if marginal taste perceptions differ among participants of normal-weight, those who are overweight and those with obesity, and whether knowledge of nutritional information affects marginal taste perception, researchers at the University of Iowa conducted a non-clinical, randomized controlled trial of 290 adults (161 with normal BMI, 78 considered overweight, and 51 considered obese) to measure instantaneous taste perceptions. Eighty percent of the participants were female, and ages ranged from 18 to 75 years. Participants were offered and rated one piece of [chocolate](#) at a time in a controlled environment and could eat as much as they wanted without feeling uncomfortable. They consumed between two and 51 pieces. Half of the study participants received nutritional information about the chocolate before the chocolate tasting began.

The study identified a consistent association between taste from food, specifically chocolate, and BMI by directly observing instantaneous taste

changes over a period of time, rather than just at the beginning and end of a period of consumption, as in prior studies.

Typically, the appeal of a specific food may decline as more of that food is eaten: the first bite of chocolate is better than the 10th, a phenomenon consistent with the concept of sensory-specific satiety. As anticipated, researchers found that ratings generally went down after each piece of chocolate consumed with no [significant difference](#) in taste perceptions between normal and overweight participants reported. However, participants with obesity had higher levels of initial taste perception, rated subsequent pieces higher than their counterparts without obesity, and their ratings declined at a more gradual rate compared to participants with normal weight and those with obesity. People hungrier prior to the study had greater taste perception; women's taste perceptions declined faster than men's; and providing nutritional information prior to chocolate consumption did not affect taste perception.

"In our study population, people with [obesity](#) reported a higher level of satisfaction for each additional piece of chocolate compared to nonobese people. Thus, their taste preferences appear markedly different," noted co-investigator Aaron C. Miller, Ph.D., Department of Epidemiology, University of Iowa, Iowa City, IA, USA. "Our findings further indicate that obese participants needed to consume a greater quantity of chocolate than nonobese participants to experience a similar decline in taste perceptions. Specifically, obese women needed to eat 12.5 pieces of chocolate to fall to the same level of taste [perception](#) as nonobese women who ate only 10 pieces, which corresponds to a difference of 67.5 calories. This may, in part, explain why obese people consume more than nonobese people."

More information: *Journal of the Academy of Nutrition and Dietetics* (2019). [DOI: 10.1016/j.jand.2019.05.018](#)

Provided by Elsevier

APA citation: Individuals with obesity get more satisfaction from their food (2019, July 30) retrieved 1

May 2021 from <https://medicalxpress.com/news/2019-07-individuals-obesity-satisfaction-food.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.