


# Intestinal immune cell interactions

1 August 2019, by Leigh Macmillan



CD8<sup>+</sup> cells promote ILC1-like intraepithelial lymphocyte homeostasis and intestinal inflammation, *PLOS ONE* (2019). [DOI: 10.1371/journal.pone.0215883](https://doi.org/10.1371/journal.pone.0215883)

Provided by Vanderbilt University

Credit: CC0 Public Domain

Multiple types of immune cells are interspersed among the epithelial cells lining the intestines and are considered to be a first line of defense against pathogens. The survival and interaction of these intraepithelial lymphocytes and their roles in intestinal inflammation are not well understood.

Ali Nazmi, Ph.D., Danyvid Olivares-Villagómez, Ph.D., and colleagues have examined the effect of one type of intraepithelial lymphocyte (iCD8 $\alpha$ ) on another (ILC1-like). Using mice with reduced numbers of iCD8 $\alpha$  cells, they found that these cells play a critical role in the survival of ILC1-like cells, and that this effect is partly mediated by the factor osteopontin.

In a mouse model of intestinal inflammation that involves ILC1-like cells, reduced iCD8 $\alpha$  cell numbers resulted in milder inflammation and treatment with osteopontin increased disease severity.

The findings, reported in *PLOS ONE*, implicate iCD8 $\alpha$  cells in ILC1-like [cell survival](#) and [intestinal inflammation](#) and add to the growing roles of osteopontin in various biological processes.

**More information:** Ali Nazmi et al. *Innate*

APA citation: Intestinal immune cell interactions (2019, August 1) retrieved 21 May 2021 from <https://medicalxpress.com/news/2019-08-intestinal-immune-cell-interactions.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*