

Transport by mobile stroke units get patients quicker treatment than ambulance

7 August 2019, by Amy Laukka



Amanda L. Jagolino-Cole, MD, and Alexandra Czap, MD, published research that highlights the advantages of hospital transport in a mobile stroke unit versus a traditional ambulance. Credit: J. Daniel Escareño/UTHealth

Every second counts for stroke patients, as studies show they can lose up to 27 million brain cells per minute. Researchers at The University of Texas Health Science Center at Houston (UTHealth) recently published new findings in *Stroke* that show patients transported to the hospital by mobile stroke unit instead of standard ambulance received a clot-busting procedure an average of 10 minutes faster, which could potentially save up to 270 million neurons per patient.

In 2014, McGovern Medical School at UTHealth was the first in the nation to launch a mobile stroke unit, a specially equipped ambulance for diagnosing and treating stroke rapidly before hospital arrival.

"The quicker we get stroke victims treatment that will restore blood flow, the more <u>brain tissue</u> we can save," said Alexandra Czap, MD, vascular neurology fellow in the Department of Neurology at

McGovern Medical School and first author of the paper. "This study shows that mobile stroke units like ours can be effective in streamlining time to treatment, potentially saving neurological function, and ultimately improving quality of life in stroke patients."

The study looked at data from 161 patients from Houston and two other locations from 2014 to 2018 who underwent intra-arterial thrombectomy after suffering an <u>acute ischemic stroke</u>, the most common kind, which is caused by a blockage in a <u>cerebral artery</u>. The only medical therapy known to treat ischemic stroke, tissue plasminogen activator (tPA), is not always able to clear a large clot. To perform the procedure, also known as endovascular thrombectomy, a vascular neurologist threads a catheter usually through a groin artery up to the blockage, where a small device inserted into the catheter is used to remove the clot.

"This is a hallmark paper because it shows that prehospital evaluation and management on a mobile stroke unit can significantly reduce time to endovascular treatment for patients with large artery clots," said Amanda L. Jagolino-Cole, MD, a teleneurologist and assistant professor in the Department of Neurology at McGovern Medical School, who was the senior author of the paper.

While on board the unit, an interdisciplinary team can begin the process to assess whether the patient needs a thrombectomy, including <u>diagnostic</u> <u>imaging</u>, neurological exam, and the administration of tPA. That gives the treatment team a <u>head start</u> and upon arrival to the hospital, the patient can be taken quickly to the endovascular suite for the procedure.

"Our mobile stroke unit allows us to bring the hospital to the patient," Czap said. "We can complete the necessary diagnostic testing and notify the hospital that we are coming, so that the appropriate teams can be ready. Streamlining this



process allowed for one of our recent mobile stroke unit patients to complete treatment in under two hours from onset of symptoms. Going forward, identification of possible thrombectomy candidates on the unit can increase the accuracy of triage and increase the number of patients having the procedure, which we hope will lead to better patient outcomes."

The paper also shows an overall improvement in endovascular <u>thrombectomy</u> metrics for all patients regardless of how they arrived to the hospital. This is significant because the procedure just started to be recognized as standard management for select <u>stroke patients</u> within the last five years, Jagolino-Cole said.

This research is part of the ongoing BEST-MSU study, a prospective comparative effectiveness trial that investigates the benefits of stroke treatment delivered using a mobile <u>stroke</u> unit compared to standard management by emergency medical services. The primary outcomes are patients' functional status at 90 days and long-term health care utilization. The BEST-MSU study will complete enrollment in 2020.

More information: Alexandra L. Czap et al, Emergency Department Door-to-Puncture Time Since 2014, *Stroke* (2019). <u>DOI:</u> <u>10.1161/STROKEAHA.119.025106</u>

Provided by University of Texas Health Science Center at Houston APA citation: Transport by mobile stroke units get patients quicker treatment than ambulance (2019, August 7) retrieved 28 April 2021 from <u>https://medicalxpress.com/news/2019-08-mobile-patients-quicker-treatment-ambulance.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.