

Osteoporosis drugs linked to reduced risk of premature death

12 August 2019

Two studies led by the Garvan Institute of Medical Research have revealed that nitrogen-bisphosphonates, drugs commonly prescribed for osteoporosis, reduced the risk of premature mortality by 34% in a cohort of over 6,000 individuals. This reduction in early mortality risk was significantly associated with a reduction in bone loss compared with no treatment.

The findings present new advice of the significant benefits of taking approved osteoporosis medicine for those at risk of osteoporosis, and their [health care](#) professional.

After the age of 50, 40% of women and 25% of men will sustain an osteoporotic fragility fracture in their life, an injury that puts them at risk of further fractures. However, currently fewer than 30% of women and 20% of men with fragility fractures are taking approved treatments for osteoporosis.

"It's a common misconception that osteoporosis affects only women, and many people choose to not take recommended treatments," says Professor Jacqueline Center, who heads the Clinical Studies and Epidemiology laboratory at the Garvan Institute and is an Endocrinologist at St Vincent's Hospital, who led the studies. "But osteoporotic fractures are not benign. Osteoporosis medication not only decreases the risk of further fractures—but it appears that this same medication also decreases [mortality rates](#) over the subsequent 15 years."

Reduction in mortality risk

Osteoporosis affects around 200 million people worldwide, and is a progressive disease in which bones become more porous and fragile, often without symptoms until the first fracture occurs.

A Garvan-led team of international researchers analysed data from a cohort of 6,120 participants aged over 50, who took part in the observational

Canadian Multicentre Osteoporosis Study.

The analysis showed that individuals treated with nitrogen-bisphosphonates (alendronate or risedronate) had a 34% reduction in mortality risk over the subsequent 15 years, compared to non-treated individuals. The study was published in the April issue of the journal *Osteoporosis International*.

In a second follow-up study, published in the *Journal of Bone and Mineral Research*, the team analysed data from a cohort of 1,735 women, from the same study. The analysis revealed that 39% of the reduction in premature mortality risk was mediated through a reduction in the rate of [bone loss](#).

The researchers also directly compared the nitrogen-bisphosphonates (alendronate or risedronate) with a weaker, non-nitrogen bisphosphonate and found a similar reduction in [mortality](#) risk benefit with the nitrogen-bisphosphonates.

The study provides additional evidence that nitrogen-bisphosphonate treatment can provide significant benefits for those with osteoporosis and is the first to examine potential mechanisms.

"For many individuals with osteoporosis, bone health isn't front-of-mind," says first author of both studies, Garvan's Dr. Dana Bliuc, Research Officer in the Clinical Studies and Epidemiology laboratory. "We hope our study results will encourage people with [osteoporosis](#) or at risk of a fracture to seek treatment—and commit to taking it."

More information: Mortality risk reduction differs according to bisphosphonate class: a 15-year observational study, *Osteoporosis International* (2019). [DOI: 10.1007/s00198-018-4806-0](https://doi.org/10.1007/s00198-018-4806-0)

Dana Bliuc et al, Reduced Bone Loss Is Associated With Reduced Mortality Risk in Subjects Exposed

to Nitrogen Bisphosphonates: A Mediation Analysis,
Journal of Bone and Mineral Research (2019). DOI:
[10.1002/jbmr.3816](https://doi.org/10.1002/jbmr.3816)

Provided by Garvan Institute of Medical Research

APA citation: Osteoporosis drugs linked to reduced risk of premature death (2019, August 12) retrieved 4
June 2022 from

<https://medicalxpress.com/news/2019-08-osteoporosis-drugs-linked-premature-death.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.