

Study of skiers holds surprises about A-fib, stroke and intense exercise

26 August 2019



Exercise is important for health and wellbeing. But past studies suggest high-intensity exercise may be a risk factor for an irregular heart rhythm called atrial fibrillation that sometimes leads to stroke. So, are athletes who develop A-Fib at higher risk for stroke?

Researchers in Sweden looked at men and women who completed one or more races in the 30- to 90-kilometer cross-country skiing event called Vasaloppet from 1989 through 2011, comparing them to a group of non-skiers. The investigators hypothesized [high-intensity exercise](#) would increase the risk of A-Fib, thus leading to more strokes in the ski group. But they discovered the opposite may be true.

"Although the group who had the best performance in a ski race had a higher risk of [atrial fibrillation](#), they didn't have a higher risk of having a stroke," said Dr. Kasper Andersen, senior author of the study published Monday in the journal *Circulation*.

Overall, male skiers had a lower risk of stroke after an A-Fib diagnosis compared to non-skiers with A-Fib, and their risk of dying from stroke was lower as well. Among women, the news was even better for skiers: They were less likely to develop A-Fib and less likely to have a stroke compared with women who didn't ski.

"Women (skiers) really didn't have an increase in atrial fibrillation, and that hasn't been seen before," said Dr. Kevin Monahan, director of the Electrophysiology and Arrhythmia Center at Boston Medical Center, who was not involved in the new study.

If A-Fib is a risk factor for stroke, why don't high-intensity athletes who develop it have a higher incidence of stroke?

According to Andersen, A-Fib is only one of the many [risk factors](#) for stroke, and [exercise](#) is one of the best ways to reduce the risk of those other factors, such as high blood pressure and diabetes.

According to American Heart Association statistics, 2.7 million to 6.1 million people in the United States have A-Fib. Risk factors for A-Fib include older age, [high blood pressure](#), diabetes and underlying heart disease. The most common symptom is a fluttering or quivering heartbeat.

The risk of developing A-Fib because of high-intensity exercise is low, Monahan said, noting the findings should not cause alarm for the average person.

"At the very extremes, for people who are training hours and hours per day, maybe there's a small risk of atrial fibrillation," he said. "It's a very small segment of the population that would be affected."

Andersen advised people with A-Fib to follow their doctor's recommended treatment to lower stroke risk because "well-trained skiers who had

developed atrial fibrillation still had a higher risk of [stroke](#) compared to both skiers and non-skiers *without* atrial [fibrillation](#)."

Both Andersen and Monahan also encourage people in general to exercise more.

"In the overall context, exercise is very good for you," Monahan said. "I think people should go out there and achieve the best level of fitness that they can."

American Heart Association News covers heart and brain health. Not all views expressed in this story reflect the official position of the American Heart Association. Copyright is owned or held by the American Heart Association, Inc., and all rights are reserved. If you have questions or comments about this story, please email editor@heart.org.

APA citation: Study of skiers holds surprises about A-fib, stroke and intense exercise (2019, August 26) retrieved 29 September 2022 from <https://medicalxpress.com/news/2019-08-skiers-a-fib-intense.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.