

## New strategy to treat Parkinson's disease

17 October 2019



Immunohistochemistry for alpha-synuclein showing positive staining (brown) of an intraneural Lewy-body in the Substantia nigra in Parkinson's disease. Credit: Wikipedia

Northwestern Medicine scientists have used patient-derived neurons to develop and test a new strategy to treat Parkinson's disease by mitigating the effects of harmful genetic mutations, as detailed in a study published today (Oct. 16) in *Science Translational Medicine.* 

Some experimental treatments for genetic disorders target mutated proteins or enzymes, but this study, led by Dr. Dimitri Krainc, took a different approach. Instead of trying to fix broken enzymes, the scientists amplified healthy ones, an approach that successfully alleviated symptoms of Parkinson's disease (PD) in human brain cells and in mouse models.

"This study highlights wild-type GCase activation as a potential therapeutic target for multiple forms of Parkinson's disease," said Krainc, who is chair of neurology and director of the Center for Neurogenetics at Northwestern University Feinberg

## School of Medicine

Parkinson's is the second-most common neurodegenerative disorder, predominately affecting <u>neurons</u> in an area of the brain called the substantia nigra. These neurons are responsible for producing dopamine—a chemical messenger used to transmit signals throughout the brain—and for relaying messages that plan and control body movement.

Mutations in the gene GBA1 represent the most common genetic risk factor for PD, according to the study, and GBA1 codes for an enzyme called glucocerebrosidase (GCase) that is important for neuronal function. PD-associated mutations can disable GBA1 and produce misshapen GCase enzymes, which contribute to an accumulation of toxic proteins in dopamine-producing neurons.

As this neuronal population dies, patients experience symptoms such as tremors and slowness of movement. While some medications can offer relief for these symptoms, there is no treatment that can stop or slow the disease.

According to Krainc, <u>drug development</u> for patients with GBA1-linked Parkinson's has largely focused on stabilizing mutated GCase and limiting its harmful effects. However, these treatments would be effective only in a few types of PD.

"Instead, activating wild-type GCase may be more relevant for multiple forms of PD that exhibit reduced activity of wild-type GCase," Krainc said.

In the current study, scientists developed a new series of chemical activators that stabilized and amplified normal GCase. The activator, a <u>small</u> <u>molecule</u> that binds to GCase, improved PD-related cellular dysfunction in patient derived neurons.

Importantly, these activators worked in several varieties of PD, showing this strategy could work for a wide range of patients, Krainc said.



"Our work points to the potential for modulating wildtype GCase activity and protein levels in both genetic and idiopathic forms of PD and highlights the importance of personalized or precision neurology in development of novel therapies," he said.

A 2017 study led by Krainc and published in Science found that some of the key pathological features of PD were only seen in human neurons and not in mouse models, further emphasizing the value of patient-derived neurons for drug development in Parkinson's disease.

"It will be important to examine human neurons to test any candidate therapeutic interventions that target midbrain dopaminergic neurons in PD," Krainc said.

**More information:** Lena F. Burbulla et al. A modulator of wild-type glucocerebrosidase improves pathogenic phenotypes in dopaminergic neuronal models of Parkinson's disease, *Science Translational Medicine* (2019). DOI: 10.1126/scitransImed.aau6870

Provided by Northwestern University APA citation: New strategy to treat Parkinson's disease (2019, October 17) retrieved 27 April 2021 from <u>https://medicalxpress.com/news/2019-10-strategy-parkinson-disease.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.