

Study finds caregiving can affect quality of life for up to a year after loss

October 30 2019



Credit: CC0 Public Domain

Family members who have been the primary carer for a dying loved one

may experience considerable grief and poor health and quality of life for several months after the person has died, new research led by Curtin University has found.

The research, published in *Palliative Medicine*, aimed to determine whether family carers' grief, general health and quality of life worsened following a loved one's death, compared to non-carers. It also examined whether the grief a carer experienced while caring could predict their grief following the death.

Lead author Associate Professor Lauren Breen, from the School of Psychology at Curtin University, said family carers are integral to providing end-of-life care for people receiving palliative care, but the tasks of caregiving can have a significant and lasting impact on them.

"Caring for a dying loved one can be a [full-time](#) and exhaustive commitment, but the grief, general health and quality of life that the carer experiences before and after death can sometimes be overlooked," Associate Professor Breen said.

"By measuring how carers were feeling in the [time period](#) leading up to the death, and then at three different time periods following the death, we were able to show that it took nine to 10 months for the carers' grief, general health, and quality of life to return to 'normal'.

"We also found that while caring, the carers' quality of life and [general health](#) was low and their grief was similar to what they experienced three to four months following the death."

Associate Professor Breen explained that the findings highlighted the need for services to provide adequate support for carers.

"Our findings present an opportunity for palliative care research and

practice to refocus efforts towards supporting carers along the caring and bereavement trajectory," Associate Professor Breen said.

"Our findings show that tailored support is needed to meet the specific needs of carers and that this should be available both before and after the death. Interventions are needed to help carers prepare for the death, improve their health and [quality of life](#), and promote adaptive grieving."

More information: Lauren J Breen et al. Effect of caregiving at end of life on grief, quality of life and general health: A prospective, longitudinal, comparative study, *Palliative Medicine* (2019). [DOI: 10.1177/0269216319880766](#)

Provided by Curtin University

Citation: Study finds caregiving can affect quality of life for up to a year after loss (2019, October 30) retrieved 8 February 2023 from <https://medicalxpress.com/news/2019-10-caregiving-affect-quality-life-year.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--