

# Improving the odds for patients with heart pumps

November 19 2019, by Jim Shelton

---



A left ventricular assist heart pump, the kind often used by patients with end-stage heart failure. Credit: Dan Renzetti

A new Yale study shows that some patients being treated for severe heart failure with a battery-operated pump saw significant improvement after additionally using neurohormonal blockade (NHB) drug therapy.

NHB therapy, which includes three broad categories of drugs, including ACE inhibitors, has long been the standard therapy for treating [heart failure](#). Until now, however, NHB therapy's effectiveness had not been extensively studied for [heart failure patients](#) using left [ventricular assist devices](#) (LVADs). The devices, which are surgically implanted, have become increasingly popular in recent years because there are few therapeutic options for end-stage patients other than heart transplants.

A new study published Nov. 18 in the journal *JAMA Cardiology* shows a clear association between the use of NHB therapy and increased survival and quality of life for patients with LVADs.

Patients who were taking any combination of the three major heart failure therapies that comprise NHB had a 56% survival rate, compared to 43.9% for patients who were not taking the medications, the researchers said.

The study, led by Nihar R. Desai, M.D., associate professor of medicine and investigator at the Yale Center for Outcomes Research & Evaluation (CORE), looked at more than 12,000 patients with LVADs in the Interagency Registry for Mechanically Assisted Circulatory Support, a North American registry, from 2008 until 2016. The analysis included patients from more than 170 medical centers in the United States and Canada.

"When we looked at the patients who were on the most intensive regimen of heart failure therapies, we found that they had one-third the risk of dying compared to those not on any of the [heart failure](#) therapies," said Megan McCullough, M.D., first author of the study.

One of the most surprising findings, McCullough and Desai said, was the inconsistent use of NHB [therapy](#) by [health care providers](#). Some providers, they noted, believe that patients with LVADs might be too

sick to benefit from the therapies.

"This study shows that the same medicines that improve quality of life and outcomes for patients without LVADs can help patients with LVADs," Desai said. "A doctor caring for an LVAD patient, after that patient is stabilized, can start to think about them as having the potential to live for years to come. There are therapies to help achieve this."

**More information:** Megan McCullough et al. Neurohormonal Blockade and Clinical Outcomes in Patients With Heart Failure Supported by Left Ventricular Assist Devices, *JAMA Cardiology* (2019). [DOI: 10.1001/jamacardio.2019.4965](https://doi.org/10.1001/jamacardio.2019.4965)

Provided by Yale University

Citation: Improving the odds for patients with heart pumps (2019, November 19) retrieved 3 January 2023 from <https://medicalxpress.com/news/2019-11-odds-patients-heart.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--