

Study examines the effects of weight loss surgery between pregnancies

29 November 2019



outcomes following bariatric surgery between a first and second pregnancy: a population data linkage study, *BJOG: An International Journal of Obstetrics & Gynaecology* (2019). DOI: [10.1111/1471-0528.15993](https://doi.org/10.1111/1471-0528.15993)

Provided by Wiley

Credit: CC0 Public Domain

During 2002–2014, there was a 13-fold increase in weight loss surgeries among women aged 15–44 years in New South Wales, Australia, and undergoing such surgery between a first and second pregnancy was associated with lower risks of hypertension, preterm birth, and other outcomes in the second pregnancy.

The findings are published in *BJOG: An International Journal of Obstetrics and Gynaecology*.

The authors of the study noted that the likelihood of adverse pregnancy outcomes among women who underwent [weight loss surgery](#) did not decrease to the level observed in the general population of women who became pregnant, however.

"Weight loss surgery for the treatment of obesity is likely to improve outcomes in a subsequent pregnancy," said lead author Ibinabo Ibiebele, of The University of Sydney Northern Clinical School.

More information: I Ibiebele et al. Perinatal

APA citation: Study examines the effects of weight loss surgery between pregnancies (2019, November 29) retrieved 25 April 2021 from <https://medicalxpress.com/news/2019-11-effects-weight-loss-surgery-pregnancies.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.