


For low back pain in older adults, treatment doesn't match guidelines

January 23 2020



Credit: CC0 Public Domain

Opioids are prescribed to nearly one-third of older adults with "new and persistent" low back pain—in most cases without trying guideline-recommended pain medications or physical therapy, according to the new research by Dan Pham Ly, MD, MPP, of Harvard University. He comments, "This study raises concerns about excessive use of low-value

and potentially harmful treatments for the common problem of LBP in older adults, with under-use of evidence-based, guideline-recommended treatments."

Many Medicare Patients Don't Receive Evidence-Based Treatments for New LBP

The study used Medicare claims data on more than 162,000 older adults with new LBP from 2011 through 2014. About 70 percent of patients were women; average age was approximately 77. None had received previous opioid treatment. The analysis included information on multiple visits for LBP over the course of a year, providing data on the timing and sequence of care.

Over half of patients (54 percent) made only one healthcare visit for LBP. That's consistent with evidence that many new episodes of LBP are self-limiting. As stated in the current American College of Physicians guidelines, most patients with LBP "improve over time regardless of treatment."

Advanced imaging studies—computed tomography (CT) or magnetic resonance imaging (MRI) scans—were used in about 15 percent of patients overall, and 29 percent of those with two or more LBP visits. In about half of cases, CT or MRI scans were performed within six weeks. That's contrary to an American Academy of Family Physicians statement that most patients don't need advanced imaging studies for initial evaluation of LBP.

Opioids were prescribed to about one-fourth of patients overall, including one-third of those with two or more LBP visits. In contrast, LBP guidelines suggest that other pain relievers—including nonsteroidal anti-inflammatory drugs (NSAIDs), such as naproxen or

ibuprofen—should be tried before opioids.

Physical therapy was prescribed to only 11 percent of patients overall and 17 percent of those with two or more visits. Evidence suggests that early use of [physical therapy](#) can avoid the use of opioids in patients with LBP.

Most patients treated with opioids had not received a prescription NSAID or physical therapy. Chronic [opioid](#) use developed in about one percent of patients overall, and nearly two percent of those with two or more visits.

Low back pain is a common reason for healthcare visits—in a given year, about ten percent of people will develop a new episode of LBP. Studies examining trends in LBP treatment found increasing use of opioids, and decreased use of NSAIDs, up to 2010. The new analysis focused on more recent patterns in evaluation and treatment of LBP in Medicaid patients, including data on repeated visits over one year.

"Many patients who develop new LBP receive guideline non-concordant care such as early advanced imaging and opioids before other modalities like PT and prescription NSAIDs," Dr. Ly writes. At least in the first half of the past decade, one-third of patients making two or more LBP visits received opioids—often without having tried other recommended treatments.

Dr. Ly calls for future studies examining barriers to guideline-recommended treatments for LBP in older adults—particularly physical therapy and NSAIDs. He also points out that pain management can be challenging in [older adults](#), highlighting the need for studies to compare the safety and effectiveness of medication options.

More information: Dan P. Ly, Evaluation and Treatment Patterns of

New Low Back Pain Episodes for Elderly Adults in the United States, 2011–2014, *Medical Care* (2020). DOI: [10.1097/MLR.0000000000001244](https://doi.org/10.1097/MLR.0000000000001244)

Provided by Wolters Kluwer Health

Citation: For low back pain in older adults, treatment doesn't match guidelines (2020, January 23) retrieved 3 July 2023 from <https://medicalxpress.com/news/2020-01-pain-older-adults-treatment-doesnt.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.