


Telemonitoring plus phone counseling lowers blood pressure among black and Hispanic stroke survivors

21 February 2020



Credit: CC0 Public Domain

Adding phone-based lifestyle counseling to home blood pressure telemonitoring is an effective strategy to improve long-term blood pressure control among minority stroke survivors with uncontrolled high blood pressure, according to late breaking science presented today at the American Stroke Association's International Stroke Conference 2020.

Uncontrolled high [blood pressure](#), or hypertension, is a major predictor of racial disparities in stroke outcomes in the United States. Improving [blood pressure control](#) among minority stroke survivors is key to secondary stroke prevention. Although home blood pressure telemonitoring and lifestyle counseling by [nurses](#) have proven effective in controlling blood pressure, this study is the first time this strategy has been tested specifically among minority stroke survivors.

Researchers randomly assigned 450 black and Hispanic stroke survivors with uncontrolled blood pressure (average age 62; 51% black; 44% women) to home blood pressure telemonitoring alone with monthly feedback to primary care

providers, or home blood pressure telemonitoring plus telephone-based counseling by nurses. The nurses counseled patients via telephone on [lifestyle behaviors](#) and strategies to improve their blood pressure and reported the blood pressure readings to the patient's doctor.

After 12 months, patients who received the home blood pressure telemonitoring plus lifestyle counseling by nurses, experienced a 14-point reduction in their systolic blood pressure, while those who received only home blood pressure telemonitoring had only a 5-point drop in their systolic blood pressure.

"The magnitude of reduction in systolic blood pressure for those who received lifestyle counseling and support was much larger than we expected," said Gbenga Ogedegbe, M.D., M.P.H., lead study author, director, Division of Health and Behavior and the Center for Healthful Behavior Change in the Department of Population Health at NYU Grossman School of Medicine in New York City. "The reduction in [systolic blood pressure](#) of 14 mmHg among these patients would be expected to translate to at least a 20% decrease in stroke deaths and 34% fewer secondary strokes. These are pretty robust findings."

Ogedegbe said, "The results suggest that hypertension management in patients at high risk for recurrent stroke should involve telephone-based [lifestyle counseling](#) by a non-physician health worker coupled with home blood pressure telemonitoring. The role of the nurse case manager could be played by the patient's pharmacist or other trained non-physician health workers."

Provided by American Heart Association

APA citation: Telemonitoring plus phone counseling lowers blood pressure among black and Hispanic stroke survivors (2020, February 21) retrieved 10 September 2022 from <https://medicalxpress.com/news/2020-02-telemonitoring-lowers-blood-pressure-black.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.