

# New study shows significant increase in weight after breast cancer

25 February 2020



Credit: CC0 Public Domain

New study findings suggest that weight gain after breast cancer is a greater problem than previously thought. The first national survey on weight after breast cancer in Australia, published in *BMC Cancer* journal, found close to two-thirds (63.7%) of women reported weight gain at an average of nine kilograms after a breast cancer diagnosis, and overall nearly one-in-five women (17%) added more than 20 kilograms.

Researchers from NICM Health Research Institute, Western Sydney University and ICON Cancer Centre, Sydney Adventist Hospital, Wahroonga, surveyed 309 [women](#) with breast cancer living in Australia using an anonymous, self-administered online cross-sectional survey between November 2017 and January 2018. The national sample consisted mainly of members from Breast Cancer Network Australia (BCNA).

The majority of women surveyed (77%) reported gaining [weight](#) within the first 12-18 months after diagnosis, which could be the ideal 'window of opportunity' to provide additional support for weight management among women with breast cancer

says Dr. Ee, lead author, [general practitioner](#) and senior research fellow at NICM Health Research Institute, Western Sydney University.

"As well as significant [weight gain](#), we also found high levels of concern about weight among our survey participants. Timing may be the key in helping women to manage weight after a diagnosis of breast cancer," said Dr. Ee.

"Cancer services and general practitioners play an important role in having early conversations with women, and referring them to a team of qualified healthcare professionals such as dietitians and exercise physiologists with experience in cancer," she said.

The survey also found:

- The proportion of women who were overweight or obese increased from 48% at time of diagnosis to 67% at the time of the survey, with the proportion of women who were obese almost doubling from 17% to 32%.
- The majority (69%) of women gained weight in excess of the rates reported in age-matched controls without breast cancer—an additional 2.42 kilograms over five years.

Professor John Boyages AM, co-author and radiation oncologist at Icon Cancer Centre says that all women should be prescribed exercise after being diagnosed with breast cancer, according to the Clinical Oncology Society of Australia (COSA) guidelines.

"As doctors we really need to actively think about weight, nutrition and exercise and advise about possible interventions," said Professor Boyages.

"Breast cancer is the most common cancer in women worldwide and in Australia, and weight gain is common after breast cancer treatment. Many

patients assume they will lose weight. Weight gain adds to self-esteem problems, increases the risk of heart disease and other cancers and several reports suggest it may affect prognosis and also increases the risk of arm swelling (lymphoedema). Prescribing a healthy lifestyle is just as important as prescribing tablets."

Dr. Ee further says that after diagnosis of breast cancer, many women experience fatigue, which can be a barrier to staying active, and studies show exercise is an effective treatment for fatigue. However, for this to be feasible and sustainable, she says supervision by an experienced exercise physiologist is invaluable.

BCNA CEO Kirsten Pilatti says many studies highlight the importance of specialists such as exercise physiologists and nutritionists for breast cancer treatment plans, however they are often an unfunded component of follow-up care.

"For many breast cancer survivors, the cost of accessing the expertise of these specialists puts them beyond their reach. We want a system that helps women and men diagnosed with [breast cancer](#) to come out and be able to move on with their life, not be crushed by the experience," said Ms Pilatti.

The researchers will next analyse the survey data to investigate reasons why women are gaining weight after [breast cancer](#), with several risk factors reported in other studies, including the type of treatment that women receive, and whether or not they were menopausal before diagnosis and treatment.

**More information:** Carolyn Ee et al, Weight before and after a diagnosis of breast cancer or ductal carcinoma in situ: a national Australian survey, *BMC Cancer* (2020). DOI: [10.1186/s12885-020-6566-4](https://doi.org/10.1186/s12885-020-6566-4)

Provided by Western Sydney University

APA citation: New study shows significant increase in weight after breast cancer (2020, February 25) retrieved 9 June 2022 from <https://medicalxpress.com/news/2020-02-significant-weight-breast-cancer.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*