


Polyps and colorectal cancer risk

17 March 2020



Risk of colorectal cancer in patients with polyps representing villous adenoma. Credit: Karolinska Institutet

Patients with polyps have a higher risk of colorectal cancer, and those with sessile serrated polyps, tubulovillous adenomas, and villous adenomas had a higher colorectal cancer mortality.

Colorectal cancer is the third most common cancer and the second leading cause of cancer death worldwide. Endoscopic screening reduces colorectal cancer and mortality by detection and removal of lesions that may develop to CRC. To prevent subsequent cancer, individuals diagnosed with either conventional adenomas or sessile serrated [polyps](#) by screening endoscopy are advised to undergo colonoscopy [surveillance](#) at different intervals, depending on the most advanced findings of the index endoscopy. However, existing guidelines for colonoscopy surveillance vary widely and lack sufficient evidence

In a matched nationwide cohort study in Sweden using individual level data from the [ESPRESSO study](#) linked to nationwide health registers for the period 1993-2016, researchers from Karolinska

Institutet, Harvard T.H. Chan School of Public Health, and Massachusetts General Hospital in the U.S examined the risk of colorectal cancer in patients with 178 377 different types of polyps and 864 831 matched reference individuals.

The 10-year cumulative incidence of colorectal cancer was 1.6% (95% CI 1.5–1.7) for hyperplastic polyps, 2.7% (2.5–2.9) for tubular adenomas, 2.5% (1.9–3.3) for sessile serrated polyps., 5.1% (4.8–5.4) for tubulovillous adenomas, and 8.6% (7.4–10.1) for villous adenomas compared with 2.1% (2.0–2.1) in reference individuals.

"The study provides detailed subtype-level data for the risk of colorectal cancer following polypectomy. The findings highlight the importance of close surveillance for patients with certain subtypes of polyps," says first author and lead investigator Mingyang Song, Massachusetts General Hospital.

Compared with reference individuals, patients with any polyps had an increased risk of colorectal [cancer](#), with multivariable HR of 1.11 (95% CI 1.02–1.22) for hyperplastic polyps, 1.77 (1.34–2.34) for sessile serrated polyps, 1.41 (1.30–1.52) for tubular adenomas, 2.56 (2.36–2.78) for tubulovillous adenomas, and 3.82 (3.07–4.76) for villous adenomas.

According to senior author and KI professor Jonas F Ludvigsson, the study suggests that follow-up of patients with different polyps could be differentiated.

"Our results indicate that patients with sessile serrated polyps, tubulovillous adenomas, and villous [adenomas](#) might benefit from surveillance, but further studies are needed to examine the impact of colonoscopy surveillance on prevention of [colorectal cancer](#)."

More information: Mingyang Song et al. Risk of colorectal cancer incidence and mortality after polypectomy: a Swedish record-linkage study, *The Lancet Gastroenterology & Hepatology* (2020).

[DOI: 10.1016/S2468-1253\(20\)30009-1](https://doi.org/10.1016/S2468-1253(20)30009-1)

Provided by Karolinska Institutet

APA citation: Polyps and colorectal cancer risk (2020, March 17) retrieved 27 April 2021 from <https://medicalxpress.com/news/2020-03-polyps-colorectal-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.