

Blocking stress-induced cocaine addiction relapse

9 April 2020, by Bill Snyder



heteroreceptors. Guanfacine should be further explored as a potential treatment for cocaine and other drug use disorders, they conclude.

More information: Rafael E. Perez et al. α 2A-adrenergic heteroreceptors are required for stress-induced reinstatement of cocaine conditioned place preference, *Neuropsychopharmacology* (2020). [DOI: 10.1038/s41386-020-0641-z](https://doi.org/10.1038/s41386-020-0641-z)

Provided by Vanderbilt University

Credit: CC0 Public Domain

Stress is a precipitating factor for craving and relapse in cocaine use disorder. A part of the brain known as the bed nucleus of the stria terminalis (BNST) has been linked to both anxiety and addiction.

Guanfacine, a drug that acts on both α 2A-adrenergic autoreceptors and heteroreceptors in the BNST, decreases stress, drug craving and [withdrawal symptoms](#) in clinical trials. But whether it can reduce relapse rates is not known.

Reporting last month in the journal *Neuropsychopharmacology*, Rafael Perez, Danny Winder, Ph.D., and colleagues demonstrate in a [mouse model](#) that heteroreceptors are required for stress-induced "reinstatement" of cocaine-seeking behavior. This finding suggests that within the BNST α 2A-adrenergic auto- and heteroreceptors may play opposing roles.

The researchers also found that low-dose guanfacine did not increase BNST activity but blocked [stress](#)-induced relapse, suggesting that at low doses, the drug does not engage

APA citation: Blocking stress-induced cocaine addiction relapse (2020, April 9) retrieved 26 July 2022 from <https://medicalxpress.com/news/2020-04-blocking-stress-induced-cocaine-addiction-relapse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.