

# How long are you infectious when you have coronavirus?

13 April 2020, by Tambri Housen, Amy Elizabeth Parry and Meru Sheel



Credit: Shutterstock

As the coronavirus pandemic stretches on, a [small proportion](#) of Australians infected have now died, while [most have either recovered](#), or are likely to recover over the next few weeks.

One thing many of us want to know is for how long people who have [SARS-CoV-2](#), the virus that causes [COVID-19](#), are able to pass it on to someone else.

Let's look at what the science tells us so far.

## How long does it take to get sick?

The "[incubation period](#)" is the time between being exposed to the virus and the onset of symptoms.

For COVID-19, the [incubation period](#) ranges from 1 to 14 days. But most people who develop COVID-19 symptoms do so [4 to 6 days](#) after exposure.

## How long are you infectious?

The "[infectious period](#)" means the time you're able to spread the virus to someone else.

For COVID-19, there is emerging evidence to

suggest the [infectious period may start](#) 1 to 3 days before you [develop symptoms](#).

The most infectious period is thought to be 1 to 3 days before symptoms start, and in the [first 7 days](#) after symptoms begin. But some people may remain infectious for longer.

Commonly reported symptoms for COVID-19—such as fever, cough and fatigue—usually last around [9 to 10](#) days but this can be longer.


## Why are some people infectious for longer?

Typically with viruses, the higher the [viral load](#) (the more virus circulating in the body), the higher the risk of transmission through known transmission pathways.

A study conducted in Hong Kong looking at viral load in 23 patients diagnosed with COVID-19 found higher viral loads in the [first week of illness](#).

Another [study from China](#) looking at 76 hospitalised patients found that by 10 days after [symptom onset](#), mild cases had cleared the virus. That is, no virus was detectable through testing.

## Coronavirus progression in majority of cases



Credit: The Conversation

However, [severe cases have much higher viral loads](#) and many continue to test positive beyond the 10 days after symptoms start.

So the more severe the illness and the higher the viral load, the longer you continue to shed the virus and are infectious.

### When are you no longer infectious?

If someone has been symptom-free for [3 days](#) and they developed their first symptoms more than [10 days](#) prior, they are no longer considered to be infectious.

But we're not sure whether people are infectious when they have recovered but the virus can still be detected in their bodies.

One study from Hong Kong found the virus could be detected for [20 days or longer](#) after the initial onset of symptoms in one-third of patients tested.

Another study from China found the virus in a

patients' faecal samples [five weeks](#) after the first [onset of symptoms](#).

But the detection of the virus doesn't necessarily mean the person is infectious. We need more studies with larger sample sizes to get to the bottom of this question.

### Should you get tested again before going back into the community?

Due to a global shortage of [coronavirus](#) tests, the [Commonwealth](#) and state governments have strict criteria about who should be tested for COVID-19 and when.

People who have been [self-quarantining](#), because they had contact with a confirmed case of COVID-19 and have completed their 14-day quarantine period without developing symptoms, can [return to the community](#). There is no requirement to be tested prior to returning to the community. It is, however, recommended they continue to practise [social distancing](#) and [good hygiene](#) as a precaution.

The requirements are different for people who have been diagnosed with COVID-19.

At present, re-testing people who have experienced mild illness, and have recovered from COVID-19 is not recommended. A person is considered safe to [return to the community](#) and discontinue self-isolation if they are no longer infectious. This means they developed their first symptoms more than [10 days](#) prior and have not experienced any symptoms for at least 3 days (72 hours).

For people who have been hospitalised with more severe illness, the [testing requirements](#) before discharge are different. They will have two swabs taken 24 hours apart to check if they have cleared the [virus](#). If the swabs are both negative, they can be discharged and don't require further self-isolation.

If one or both tests are positive but the person is well enough to go home, they must continue to self-isolate for at least 10 days since they were discharged from hospital and they have not

experienced any symptoms for at least 3 days.

There are also different testing requirements for people working or living in [high-risk settings](#). If you work or live in a high-risk setting you should consult with your health care provider on re-testing requirements.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

APA citation: How long are you infectious when you have coronavirus? (2020, April 13) retrieved 19 October 2022 from <https://medicalxpress.com/news/2020-04-infectious-coronavirus.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*