


# Age, sex and smoking influence opioid receptor function in the brain

May 28 2020



Distribution of  $\mu$ -opioid receptors in the human brain. Warmer colour stands for more receptors, and colder colour for less receptors. Credit: *NeuroImage*

Opioids regulate the feelings of pleasure and pain in the brain. A study by the national Turku PET Centre in Finland shows that age, sex and smoking influence  $\mu$ -opioid receptor density in the brain. The results of the study help to better understand the differences between individuals when it comes to neuropsychiatric disorders.

Dysfunctions of the brain's opioid system are associated with several

disorders, such as addiction, and chronic pain problems.

"We noticed in our study that age, sex and smoking have fundamental influence on the organisation of the human brain's opioid system. This suggests that there are significant differences in the opioid system between individuals, which may explain why some individuals are prone to develop opioid-linked pathological states, such as psychiatric disorders," explains Tatu Kantonen, Doctor of Medicine.

Brain's  $\mu$ -opioid receptors act as important mediators for body's own opioids in the brain. This study analysed [positron emission tomography](#) (PET) brain scans targeting  $\mu$ -opioid receptors from 204 individuals with no neurologic or [psychiatric disorders](#). This database was compiled with new computational tools developed at the Turku PET Centre.

"When an opioid molecule binds to a dock-like receptor in the brain, it may cause a feeling of reduced pain or increased pleasure. Putting it more simply, opioids are like the body's own ambassadors of pleasure and relief," explains Kantonen.

Older age was associated with increased  $\mu$ -opioid receptors in the [cerebral cortex](#), whereas in the deeper parts of the brain they were decreased. The increase in receptors was stronger in males. Smoking was associated with decreased  $\mu$ -opioid receptors in most [brain regions](#).

The researchers also discovered that there are more  $\mu$ -opioid receptors in the right versus the left hemisphere of the brain. This observation may help to explain the previously reported differences between the two hemispheres in the processing of emotions and pain.

**More information:** Tatu Kantonen et al. Interindividual variability and lateralization of  $\mu$ -opioid receptors in the human brain, *NeuroImage* (2020). [DOI: 10.1016/j.neuroimage.2020.116922](https://doi.org/10.1016/j.neuroimage.2020.116922)

Provided by University of Turku

Citation: Age, sex and smoking influence opioid receptor function in the brain (2020, May 28)  
retrieved 10 July 2023 from <https://medicalxpress.com/news/2020-05-age-sex-opioid-receptor-function.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.