


For each day's delay in social distancing, a COVID-19 outbreak lasts days longer

1 June 2020, by Esther Robards-Forbes



Credit: CC0 Public Domain

A new analysis of COVID-19 outbreaks in 58 cities has found that places that took longer to begin implementing social distancing measures spent more time with the virus rapidly spreading than others that acted more quickly.

In a new paper from epidemiological researchers at The University of Texas at Austin, now in press with the CDC's journal *Emerging Infectious Diseases*, researchers studied cities throughout China and analyzed when first cases were detected, when social distancing measures were implemented and when the [outbreak](#) was considered contained. The team found that every day a [city](#) delayed in implementing social distancing measures after the appearance of a first case added 2.4 days to the length of the outbreak.

"Every day saves time, saves effort, saves people becoming infected and probably saves lives," said Lauren Ancel Meyers, a professor of integrative biology who leads the UT Austin COVID-19 Modeling Consortium. "This is particularly important as we think about the coming weeks and months."

The findings are applicable both to communities experiencing their first outbreaks and those that

may see a resurgence in the coming months, Meyers said. Waiting a week after early signs of resurgence might require about 17 more days of social distancing to slow the spread of the pandemic, according to the data.

"It will be difficult to consider strict interventions again, but acting early upon signs of resurgence will mean fewer days of social distancing orders," said Spencer Fox, associate director of the UT Austin COVID-19 Modeling Consortium and co-author on the paper.

"Our findings have implications for the timing of interventions in U.S. cities," Meyers said. "The impact of delays may be particularly important for communities that are prone to rapid transmission, such as nursing homes, colleges, schools and jails. We need concrete plans for when and how to respond to rising cases to prevent unnecessarily long and costly restrictions."

To determine when an outbreak had been contained, scientists look at case counts and determine the reproduction number, a measurement that shows how many people will be infected by one infectious person. If the reproduction number drops below 1, scientists consider the outbreak contained.

Other studies into the impact of delaying social distancing measures have used modeling to estimate a link between the time of measures being taken and the effect on outbreaks. By contrast, this study used on-the-ground data to determine the link between cities taking measures to contain the virus and rates of the virus spreading.

The study was not able to determine which social distancing measures were most effective, but it showed that the timing of the first measure, regardless of the type, had a big impact.

"We provide direct, data-driven evidence that the

timing of interventions has a substantial impact on how long an outbreak lasts, how effective our interventions are and, ultimately, how many people might be infected and die from the virus," Meyers said.

Although the study looked at cities experiencing the earliest days of an outbreak, the findings are also relevant for cities in the middle of an outbreak, Meyers said.

Provided by University of Texas at Austin

APA citation: For each day's delay in social distancing, a COVID-19 outbreak lasts days longer (2020, June 1) retrieved 15 October 2022 from <https://medicalxpress.com/news/2020-06-day-social-distancing-covid-outbreak.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.