

Scientists reveal involvement of dorsal periacqueductal gray in aversive conditioning

11 June 2020, by Li Yuan



Credit: CC0 Public Domain

The dorsal periacqueductal gray (dPAG) is involved in controlling emotional behavior by coordinating defensive behavior. It is linked to pathologies such as post-traumatic disorder (PTSD). However, its role in the formation and retrieval of negative memories remains unknown.

Researchers from the Shenzhen Institutes of Advanced Technology (SIAT) of the Chinese Academy of Sciences have investigated the role and connectivity of the [glutamatergic](#) (VGluT2+) and GABAergic (GAD2+) subpopulations of neurons in the dPAG of mice, in the context of aversive conditioning and retrieval.

By recording activity using in vivo and free-moving calcium imaging, the researchers found that VGluT2+ and GAD2+ subpopulations similarly responded to fear unconditioned and conditioned stimulations.

But 24 hours after conditioning, only the VGluT2+

subpopulation persisted in responding to the conditioned stimulation during retrieval test.

These results indicate that glutamatergic dPAG subpopulation was involved in aversive [memory](#), but not GABAergic dPAG subpopulation.

This intrigued SIAT team, since it indicated functional differences between glutamatergic and GABAergic populations during the integration of aversive memories.

To better understand these divergences, they mapped the structures projecting respectively to dPAG^{VGluT2+} and dPAG^{GAD2+} neurons by using Rabies Virus, a Cre-dependent monosynaptic retrograde tracing technique.

The results showed that both glutamatergic and GABAergic dPAG subpopulations received upstream projections from the same structures, but with different strength of inputs. In particular, compared with dPAG^{VGluT2+}, the dPAG^{GAD2+} population received many more Ventromedial Hypothalamus inputs, a structure known to be involved in sustained fear emotional states.

These specific patterns of connectivities could explain part of the functional results obtained, and may especially indicate that glutamatergic subpopulation is a main contributor of aversive memories in dPAG.

"Investigating how structures belonging to emotion circuitry, such as the PAG, are involved in aversive memory formation is an important step to understand long-lasting emotional states. Our study could ultimately lead to find potential treatments for pathologies such as PTSD," said Dr. Quentin Montardy, the paper's first co-author.

The study was published in *Neuroscience Letters* on May 23, 2020.

More information: Quentin Montardy et al.
Glutamatergic and gabaergic neuronal populations in the dorsal Periaqueductal Gray have different functional roles in aversive conditioning, *Neuroscience Letters* (2020). DOI: [10.1016/j.neulet.2020.135059](https://doi.org/10.1016/j.neulet.2020.135059)

Provided by Chinese Academy of Sciences
APA citation: Scientists reveal involvement of dorsal periaqueductal gray in aversive conditioning (2020, June 11) retrieved 1 August 2022 from <https://medicalxpress.com/news/2020-06-scientists-reveal-involvement-dorsal-periaqueductal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.