

There may be 6 types of COVID-19

July 28 2020



COVID-19 may not be just one disease, but six distinct types, a new British study claims.

Each type differs in severity and in the need for respiratory support during hospitalization, the researchers added.

Cough, fever and [loss of smell](#) are the usual symptoms of COVID-19, but the range of symptoms can include headaches, muscle [pain](#), fatigue, diarrhea, confusion, loss of appetite, shortness of breath and more.

For the study, the researchers analyzed data on 1,600 people who reported their symptoms into an app.

The six [symptom](#) groups in a sequence from least to most severe are:

1. Headache, loss of smell, muscle pains, cough, [sore throat](#), [chest pain](#), no fever.
2. Headache, loss of smell, cough, sore throat, hoarseness, fever, loss of appetite.
3. Headache, loss of smell, loss of appetite, diarrhea, sore throat, chest pain, no cough.
4. Headache, loss of smell, cough, fever, hoarseness, chest pain, fatigue.
5. Headache, loss of smell, loss of appetite, cough, fever, hoarseness, sore throat, chest pain, fatigue, confusion, muscle pain.
6. Headache, loss of smell, loss of appetite, [cough](#), fever, hoarseness, sore throat, chest pain, fatigue, confusion, muscle pain, shortness of breath, diarrhea, abdominal pain.

The last three types are tied with the most severe disease, the researchers noted. The range of those with [severe symptoms](#) who need help breathing range from about 9% to 20%, while those with milder symptoms who need breathing aids range from 2% to 3%.

Also, nearly half of the patients with the most severe symptoms ended up in a hospital, compared with 16% of those with the least severe symptoms, the findings showed.

Using a mix of symptoms, [body weight](#) and other factors, the researchers developed a model that predicts which patients will need to be hospitalized and need breathing support.

"These findings have important implications for care and monitoring of people who are most vulnerable to severe COVID-19," said researcher Dr. Claire Steves, from King's College London.

"If you can predict who these people are at day five, you have time to give them support and early interventions such as monitoring blood oxygen and sugar levels, and ensuring they are properly hydrated—simple care that could be given at home, preventing hospitalizations and saving lives," Steves said in a college news release.

The report was published online July 28 on *medRxiv*, a preprint server that does not include the rigors of peer review.

More information: For more on COVID-19, head to the [U.S. Centers for Disease Control and Prevention](#).

Copyright © 2020 [HealthDay](#). All rights reserved.

Citation: There may be 6 types of COVID-19 (2020, July 28) retrieved 23 March 2023 from <https://medicalxpress.com/news/2020-07-covid--1.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--