


Stress and anger may exacerbate heart failure

12 August 2020



Credit: David Shankbone/Wikipedia

Mental stress and anger may have clinical implications for patients with heart failure according to a new report published in the *Journal of Cardiac Failure*.

Heart failure is a life-threatening [cardiovascular disease](#) in which the heart is damaged or weakened. This can lead to a reduced ejection fraction, in which the [heart muscle](#) pumps out a lower amount of blood than is typical with each contraction.

In this study of patients who had [heart failure](#) with reduced ejection fraction, the authors—including researchers at Yale—evaluated the [effects of stress](#) and anger on diastolic function. Diastolic function describes the ability of the heart to relax and refill between muscle contractions and is predictive of mortality risk.

For one week, participants completed daily questionnaires about their experiences of stress, anger, and negative emotions during the previous 24 hours. Participants then completed a standardized "mental stress" protocol in which they solved challenging arithmetic problems and described a recent stressful experience. Echocardiograms were performed to assess diastolic function at rest and during the stress task.

Patients who reported experiencing anger in the week prior to the laboratory mental stress protocol exhibited worse baseline resting diastolic pressure, the researchers said. Furthermore, most patients demonstrated stress-provoked changes in diastolic function, including decreased early relaxation and increased diastolic pressure.

"Mental stress is common in patients with heart failure due in part to the complexities of disease self-management, progressively worsening functional limitations, and frequent symptom exacerbations and hospitalizations," said the lead author Kristie Harris, a postdoctoral associate in cardiovascular medicine at Yale.

"We have evidence that patients who experience chronically elevated levels of stress experience a more burdensome disease course with diminished quality of life and increased risk for adverse events. Clarifying the relevant behavioral and physiological pathways is especially important in the era of COVID-19 when the typical stressors of heart failure may be further compounded by pandemic-related stressors," Harris said.

"Factors such as [mental stress](#) and anger often go unrecognized and are under-addressed," said Matthew Burg, a Yale [clinical psychologist](#) and senior author of the study. "This study contributes to the extensive literature showing that stress and anger affect clinical outcomes for patients with heart disease, adding chronic heart failure to the list that includes [ischemic heart disease](#) (narrowed

arteries) and arrhythmic disease."

Burg said that while stress management and related techniques have been shown to reduce risk for adverse events among patients with ischemic heart disease (narrowed arteries), further work is needed to identify factors that increase vulnerability to the effects of stress in heart failure, and to determine whether stress management can improve outcomes for

More information: Kristie M. Harris et al, Impact of Mental Stress and Anger on Indices of Diastolic Function in Patients with Heart Failure, *Journal of Cardiac Failure* (2020). DOI: [10.1016/j.cardfail.2020.07.008](https://doi.org/10.1016/j.cardfail.2020.07.008)

Provided by Yale University

APA citation: Stress and anger may exacerbate heart failure (2020, August 12) retrieved 5 June 2022 from <https://medicalxpress.com/news/2020-08-stress-anger-exacerbate-heart-failure.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.