


Using personal frequency to control brain activity

17 August 2020



Credit: Pixabay/CC0 Public Domain

Stroke, Parkinson's disease and depression—these medical illnesses have one thing in common: they are caused by changes in brain functions. For a long time, research has therefore been conducted into ways of influencing individual brain functions without surgery in order to compensate for these conditions.

Scientists at the Max Planck Institute for Human Cognitive and Brain Sciences in Leipzig, Germany, have taken a decisive step. They have succeeded in precisely influencing the functioning of a single area of the brain. For a few minutes, they inhibited exactly the area that processes the sense of touch by specifically intervening in its rhythm. As a result, the area that was less networked with other [brain regions](#), its so-called functional connectivity, decreased, and thus also the exchange of information with other brain networks.

This was possible because the researchers had previously determined each participant's individual brain rhythm that occurs when perceiving touch. With the personal frequency, they were able to modulate the targeted areas of the brain one at a time in a very precise manner using what is known

as transcranial alternating current [stimulation](#). "This is an enormous advance," explains Christopher Gundlach, first author of the underlying study. "In previous studies, connectivity fluctuated extensively when the current was distributed in different areas of the brain. The electrical current randomly sought its own path in the brain and thus affected different brain areas simultaneously in a rather imprecise manner.

In a preliminary study, the neuroscientists had already observed that this form of stimulation not only reduces the exchange of the targeted brain networks with other networks, it also affects the brain's ability to process information, in this case the sense of touch. When the researchers inhibited the responsible somatosensory [network](#), the perception threshold increased. The study participants only perceived stimuli when they were correspondingly strong. When, on the other hand, they stimulated the region, the [threshold value](#) dropped and the study participants already felt very gentle electrical stimuli.

"The deliberate change in brain rhythm lasted only briefly. As soon as the stimulation is switched off, the effect disappears again," explains study leader Bernhard Sehm. "Nevertheless, the results are an important step towards a targeted therapy for diseases or disorders caused by disturbed brain functions." Targeted [brain](#) stimulation could help to improve, direct and, if necessary, attenuate the flow of information.

More information: Christopher Gundlach et al, Reduction of somatosensory functional connectivity by transcranial alternating current stimulation at endogenous mu-frequency, *NeuroImage* (2020). [DOI: 10.1016/j.neuroimage.2020.117175](https://doi.org/10.1016/j.neuroimage.2020.117175)

Provided by Max Planck Society

APA citation: Using personal frequency to control brain activity (2020, August 17) retrieved 28 August 2022 from <https://medicalxpress.com/news/2020-08-personal-frequency-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.