


No benefit for post-operative radiotherapy in non-small-cell lung cancer

20 September 2020



Credit: CC0 Public Domain

Post-operative radiotherapy (PORT) used in patients with non-small-cell lung cancer (NSCLC) following complete resection and after (neo) adjuvant chemotherapy shows no statistically significant difference in 3-year disease-free survival (DFS), according to data presented at ESMO 2020. These results give the oncology community a long-awaited answer.

PORT in completely resected NSCLC [patients](#) has been a subject of debate for many years in patients with mediastinal nodal involvement (pN2), since a meta-analysis in 1998 threw doubt on the benefits associated with it.

However, more recent times have seen better selection, (neo)-adjuvant chemotherapy in stage III resected patients, as well as improved radiotherapy and more recent non prospective studies suggested modern PORT could improve outcome.

As such, there was a clear need for a large, randomised trial to assess the role of modern mediastinal PORT in adequately staged and

surgically treated patients. This study provides more robust data to help clinicians to decide the best course of action for these patients.

The large randomised controlled trial presented at ESMO 2020, explored the role of modern mediastinal PORT in patients with completely resected NSCLC with histo/cytologically proven nodal involvement.

A total of 501 patients were entered into the intention-to-treat analysis, of which 252 received PORT over five weeks, and 249 entered the control arm (no PORT). Safety analysis was carried out in 487 patients.

Disease-free survival was of 47.1% in the PORT arm and 43.8% in the control arm, thus not statistically significant, with a hazard ratio of = 0.85 (95% CI = [0.67;1.07]; p value = 0.16) for patient receiving PORT compared to control.

Overall survival at three years was 66.5% (95% CI = [59;73]) of patients in the PORT arm compared to 68.5% (95% CI = [61;75]) in the control arm.

Study author Dr. Cecile Le Pechoux, radiation oncologist from Institut Gustave Roussy, Paris, France, said: "PORT cannot be recommended for all patients with stage II and III NSCLC with mediastinal nodal involvement. Possibly, however, for some patients it might be useful because it does decrease the rate of mediastinal relapse by 50%. This must be put into balance with the risk of over-added cardio-pulmonary toxicity. We need to do further analysis to determine if certain patients, in particular, could benefit from it" she added.

Prof Rafal Dziadziuszko, radiation oncologist from the Medical University of Gdansk, Poland, commented on the findings. "Radiotherapy to the mediastinum after surgery, after [adjuvant chemotherapy](#) shouldn't be recommended as standard of care. This will change the practice of

many institutions that adopted standard use of radiotherapy in these patients. We can safely say there is no net benefit from such treatment but there is also potential harm, which we see from this study, so any potential benefits in some patients are offset by the predominantly higher risk of cardiopulmonary toxicities."

More information: Abstract LBA3_PR 'An international randomized trial, comparing post-operative conformal radiotherapy (PORT) to no PORT, in patients with completely resected non-small cell lung cancer (NSCLC) and mediastinal N2 involvement. Primary end-point analysis of Lung ART (IFCT-0503, UK NCRI, SAKK) NCT00410683.' will be presented by Cecile Le Pechoux during the Presidential Symposium II, on Sunday, 20 September 2020. *Annals of Oncology*, Volume 31 Supplement 4, September 2020

Provided by European Society for Medical Oncology

APA citation: No benefit for post-operative radiotherapy in non-small-cell lung cancer (2020, September 20) retrieved 15 April 2021 from <https://medicalxpress.com/news/2020-09-benefit-post-operative-radiotherapy-non-small-cell-lung.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.