


# Lung cancer screening a step closer to reality following combined study

29 September 2020



Lung CA seen on CXR. Credit: James Heilman, MD/Wikipedia

Newly released study results present a strong case for lung cancer screening in New Zealand—particularly for Māori whose mortality rates are between three and four times higher than other ethnic groups.

Around 450 Māori are diagnosed with lung cancer each year and approximately 300 die from it.

Waitematā and Auckland DHBs have collaborated with University of Otago researchers on a study to assess whether low dose computerised tomography (LDCT) might be a cost-effective way to catch the disease in its early stages and improve [health outcomes](#) for Māori and, ultimately, other [ethnic groups](#).

LDCT is a computerised X-ray that produces very

clear three dimensional images by using only a small amount of radiation.

The study, led by Associate Professor Sue Crengle (Kōi Tahu, Kōti Mamoe, Waitaha) from the University of Otago and just published online by medical journal *BMJ Open*, used scientific modelling to estimate the benefits and costs of LDCT screening in a high-risk population.

Multiple factors need to be considered before any kind of screening programme goes ahead but an answer to the question of cost-effectiveness is a significant step forward.

"Having clarity about the cost-effectiveness of [lung cancer screening](#) provides further impetus to get this work started," Associate Professor Crengle says. "If this works for Māori, then it will work for everyone else as well."

Study findings show that:

- a national biennial lung cancer screening programme is likely to be cost-effective for Māori and the total population in New Zealand.
- screening is likely to reduce population inequities relating to lung cancer among Māori.

Screening trials have been completed overseas where they have resulted in a 20-26 percent reduction in lung cancer deaths.

The DHBs and University of Otago are now planning to trial an invitation process with up to 50 general practices across the Auckland and Waitematā catchments. The trial will be aimed at Māori patients and the results will help shape any future screening roll-out across New Zealand.

Waitematā DHB and Auckland DHB Director of Health Outcomes Dr. Karen Bartholomew says

deaths from lung cancer are the single-greatest contributor to the higher mortality rate for Māori compared with NZ Europeans / other ethnicities.

"Around 75 percent of early-stage [lung cancer](#) may be curable." Dr. Bartholomew says. "If [lung cancer](#) is caught early enough, the impact of screening could go a long way towards helping us save lives and eliminate the unacceptable inequities we currently see in mortality between Māori and non-Māori."

**More information:** Melissa McLeod et al, Impact of low-dose CT screening for lung cancer on ethnic health inequities in New Zealand: a cost-effectiveness analysis, *BMJ Open* (2020). [DOI: 10.1136/bmjopen-2020-037145](#)

Provided by University of Otago

APA citation: Lung cancer screening a step closer to reality following combined study (2020, September 29) retrieved 31 May 2022 from <https://medicalxpress.com/news/2020-09-lung-cancer-screening-closer-reality.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*