

# Research could change how blood pressure is managed in spinal cord injury patients

15 October 2020



Credit: CC0 Public Domain

New research from the International Collaboration on Repair Discoveries (ICORD) challenges the current standard for managing blood pressure in people with spinal cord injury (SCI).

The findings, published today in *Nature Communications*, could lead to a change in the way newly injured patients have their blood pressure managed, potentially improving their chances of retaining more function in the long term.

This study, led by ICORD principal investigators Dr. Christopher West and Dr. Brian Kwon, demonstrated that following high-thoracic SCI, the heart's ability to contract is impaired, leading to reduced [spinal cord](#) blood flow. High-thoracic SCI generally refers to injuries that affect the abdominal and lower back muscles and the legs, typically resulting in paraplegia, while arm and hand function may not be affected.

Currently, a patient being treated for acute SCI—a traumatic [injury](#) that bruises, partially or fully tears the spinal cord—has their blood pressure managed using drugs that cause their [blood vessels](#) to

constrict in order to increase blood pressure.

ICORD principal investigators Dr. Christopher West and Dr. Brian Kwon and their research teams have published a study that challenges the current standard for managing blood pressure in people with spinal cord injury (SCI).

The findings could lead to a change in the way newly injured patients have their blood pressure managed, potentially improving their chances of retaining more function in the long term.

This study, published today in *Nature Communications*, demonstrated that following high-thoracic SCI, the heart's ability to contract is impaired, leading to reduced spinal cord blood flow. Currently, a patient being treated for acute SCI has their blood pressure managed using drugs that cause their blood vessels to constrict in order to increase blood pressure. In this study, Drs. West and Kwon and their team used an experimental treatment targeting the heart to beat more powerfully, which increased the amount of blood ejected and also increased blood pressure.

"The key difference is that by targeting the heart, we increase [blood pressure](#) by increasing blood flow instead of by causing the blood vessels to narrow," said Dr. West. "This is important because by increasing blood flow we can deliver more blood and oxygen to the spinal cord which is expected to minimize the damage at the epicenter of the injury, and therefore reduce injury severity and bleeding in the cord, both of which are expected to improve the chances of people with SCI retaining more function."

This could mean in the future that an individual who is newly admitted to the hospital may receive a different drug during their initial period of post-injury management. If proven to be effective in humans with SCI, it could improve their chances of retaining more function in the long-term.

"It is important to note that one of the only things we can currently do for acute SCI patients is to try to optimize the supply of [blood](#) and oxygen to the injured spinal cord to prevent further secondary injury," said Dr. Kwon. "We need to seek ways of delivering this care in the most effective manner possible, as any bit of spinal cord function that could be improved upon by novel approaches such as this would potentially be very impactful to a patient."

**More information:** Alexandra M. Williams et al, Cardio-centric hemodynamic management improves spinal cord oxygenation and mitigates hemorrhage in acute spinal cord injury, *Nature Communications* (2020). [DOI: 10.1038/s41467-020-18905-8](#)

Provided by University of British Columbia

APA citation: Research could change how blood pressure is managed in spinal cord injury patients (2020, October 15) retrieved 5 May 2021 from <https://medicalxpress.com/news/2020-10-blood-pressure-spinal-cord-injury.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*