


Questionnaire-based tool measures fatigue in patients receiving dialysis

23 October 2020



Credit: Pixabay/CC0 Public Domain

A recent study has validated a new patient-reported outcome measure that assesses fatigue in patients receiving long-term dialysis treatments. The results appear in an upcoming issue of *CJASN*.

Fatigue, which affects most patients with [kidney failure](#) who receive hemodialysis, contributes to poor quality of life and is associated with higher risks of cardiovascular disease, depression, and early death. Unfortunately, fatigue can be difficult to quantify. To address this, Angela Ju, Ph.D. (University of Sydney, in Australia) and her colleagues developed a patient-reported measure of fatigue— called Standardized Outcomes in Nephrology-Hemodialysis (SONG-HD) Fatigue—and evaluated its potential in 485 adults receiving hemodialysis in the United Kingdom, Australia, and Romania. SONG-HD Fatigue calculates the average fatigue over a period of 1 week, and it gauges tiredness, energy, and the impact of fatigue on life participation.

"Currently, fatigue is assessed in a wide range of different ways using measures that vary in terms of content, length, reliability, and validity. Most are

quite lengthy, and it is unclear which measure is the best to use," said Dr. Ju. "This makes it difficult for researchers to choose how to assess fatigue, and compare effectiveness of interventions evaluated in different studies."

The study assessed SONG-HD Fatigue in accordance with the recommended US Food and Drug Administration guidelines for patient-reported outcome measures. The authors concluded that it appears to be a reliable and valid measure to be used in trials involving patients receiving hemodialysis.

"Establishing a core outcome measure for fatigue will allow researchers to do 2 important things: to quickly and accurately assess the impact a new intervention has on fatigue as experienced by patients receiving hemodialysis, and to allow the comparison of interventions across different trials to see which one is superior in terms of its effect on fatigue," Dr. Ju said. "Patients want to know about fatigue regardless of whether or not the [intervention](#) was designed to affect it. The core outcome measure helps to provide this information. Interventions specifically targeting fatigue will require a more comprehensive measure."

An accompanying editorial notes that "patients on maintenance hemodialysis have been very clear that [fatigue](#) is an important problem for them. Investigators such as Ju and colleagues have heard them. Now it is time for all us, clinicians and researchers alike, to do the same."

A Patient Voice also accompanies the study and provides the perspective of a psychologist who has late-stage chronic kidney disease.

More information: Angela Ju et al, Validation of a Core Patient-Reported Outcome Measure for Fatigue in Patients Receiving Hemodialysis, *Clinical Journal of the American Society of Nephrology* (2020). [DOI: 10.2215/CJN.05880420](https://doi.org/10.2215/CJN.05880420)

Sarah J. Ramer et al. Moving the Science of Patient-Reported Outcome Measures Forward, *Clinical Journal of the American Society of Nephrology* (2020). DOI: [10.2215/CJN.14900920](https://doi.org/10.2215/CJN.14900920)

Henning Sondergaard. Fatigue While Undergoing Long-Term Hemodialysis, *Clinical Journal of the American Society of Nephrology* (2020). DOI: [10.2215/CJN.14870920](https://doi.org/10.2215/CJN.14870920)

Provided by American Society of Nephrology

APA citation: Questionnaire-based tool measures fatigue in patients receiving dialysis (2020, October 23) retrieved 24 April 2021 from <https://medicalxpress.com/news/2020-10-questionnaire-based-tool-fatigue-patients-dialysis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.