

# Your favorite music can send your brain into a pleasure overload

November 3 2020

---



Credit: CC0 Public Domain

We all know that moment when we're in the car, at a concert or even sitting on our sofa and one of our favorite songs is played. It's the one that has that *really* good chord in it, flooding your system with pleasurable emotions, joyful memories, making your hair stand on edge, and even sending a shiver or "chill" down your spine. About half of

people get chills when listening to music. Neuroscientists based in France have now used EEG to link chills to multiple brain regions involved in activating reward and pleasure systems. The results are published in *Frontiers in Neuroscience*.

Thibault Chabin and colleagues at the Université de Bourgogne Franche-Comté in Besançon EEG-scanned the brains of 18 French participants who regularly experience chills when listening to their favorite musical pieces. In a questionnaire, they were asked to indicate when they experienced chills, and rate their degree of pleasure from them.

"Participants of our study were able to precisely indicate "chill-producing" moments in the songs, but most musical chills occurred in many parts of the extracts and not only in the predicted moments," says Chabin.

When the participants experienced a chill, Chabin saw specific electrical activity in the orbitofrontal cortex (a region involved in emotional processing), the supplementary motor area (a mid-[brain](#) region involved in movement control) and the right temporal lobe (a region on the right side of the brain involved in auditory processing and musical appreciation). These regions work together to process music, trigger the brain's reward systems, and release dopamine—a "feel-good" hormone and neurotransmitter. Combined with the pleasurable anticipation of your favorite part of the song, this produces the tingly chill you experience—a physiological response thought to indicate greater cortical connectivity.

"The fact that we can measure this phenomenon with EEG brings opportunities for study in other contexts, in scenarios that are more natural and within groups," Chabin comments. "This represents a good perspective for musical emotion research."

EEG is a non-invasive, highly accurate technique that scans for electrical currents caused by brain activity using sensors placed across the surface of the scalp. When experiencing musical chills, low frequency electrical signals called "theta activity"—a type of activity associated with successful memory performance in the context of high rewards and musical appreciation—either increase or decrease in the [brain regions](#) that are involved in musical processing.

"Contrary to heavy neuroimaging techniques such as PET scan or fMRI, classic EEG can be transported outside of the lab into naturalistic scenarios," says Chabin. "What is most intriguing is that music seems to have no biological benefit to us. However, the implication of dopamine and of the reward system in processing of musical pleasure suggests an ancestral function for music."

This ancestral function may lie in the period of time we spend in anticipation of the "chill-inducing" part of the music. As we wait, our brains are busy predicting the future and release dopamine. Evolutionarily speaking, being able to predict what will happen next is essential for survival.

Why should we continue to study chills?

"We want to measure how cerebral and physiological activities of multiple participants are coupled in natural, social musical settings," Chabin says. "Musical pleasure is a very interesting phenomenon that deserves to be investigated further, in order to understand why music is rewarding and unlock why [music](#) is essential in human lives."

How the study was done:

The study was carried out on 18 healthy participants—11 female and 7 male. Participants were recruited through posters on the campus and

university hospital. They had a mean age of 40 years, were sensitive to musical reward, and frequently experienced chills. They had a range of musical abilities.

A high-density EEG scan was conducted as participants listened to 15 minutes of 90 s excerpts of their most enjoyable musical pieces. While listening, participants were told to rate their subjectively felt pleasure and indicate when they felt "chills". In total, 305 chills were reported, each lasting, on average, 8.75 s. These findings implied increased [brain activity](#) in regions previously linked to musical pleasure in PET and fMRI studies.

**More information:** *Frontiers in Neuroscience* (2020). [DOI: 10.3389/fnins.2020.565815](#) , [www.frontiersin.org/articles/10.3389/fnins.2020.565815/full](#)

Provided by Frontiers

Citation: Your favorite music can send your brain into a pleasure overload (2020, November 3) retrieved 31 January 2023 from <https://medicalxpress.com/news/2020-11-favorite-music-brain-pleasure-overload.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--