

The hidden reason children born by C-section are more likely to develop asthma

11 November 2020



Credit: CC0 Public Domain

Researchers at Rutgers University, the Copenhagen Prospective Studies on Asthma in Childhood and the University of Copenhagen have described for the first time how delivery by caesarean section interferes with a baby's ability to obtain beneficial germs from the mother's microbiome, and how this can lead to early childhood asthma.

The study, published in *Science Translational Medicine*, has implications for understanding the role of C-section delivery in potentially skewing a child's microbiota—the trillions of beneficial microorganisms that live in and on our bodies—and how this can influence health. It could lead to new prevention strategies, including targeted efforts to fix disturbances in a child's [microbiome](#).

"Every generation of mothers hands over its microbiome to the next, as the baby is coated with

beneficial germs while being squeezed through the [birth canal](#)—but this doesn't happen for babies born through C-section," said co-author Martin Blaser, director of the Center for Advanced Biotechnology and Medicine at Rutgers. "It takes a while for babies born through C-section to develop a normal microbiome. And during that time, while the [immune system](#) is also developing, they become more at risk for later developing certain diseases like asthma. This study provides a mechanism for the known link between C-section birth and heightened risk of asthma."

The researchers analysed the effects of vaginal birth versus C-section during the first year of life for 700 children. To determine whether the type of delivery caused disruptions in the microbiome, and whether these were associated with asthma, they examined children's fecal samples at one week, one month and one year to determine microbial diversity and maturity.

The researchers found that delivery by caesarean section was associated with more than a doubled risk of later asthma and allergies, as well as significant changes in the composition of the gut microbiota. However, at one year of age, the risk of asthma was reduced in C-section-born children if their gut microbiota had recovered from its initial disruption and begun to mature normally.

"Even though a child is born by caesarean section and has an immense early microbial perturbation, this may not lead to a higher risk of asthma if the microbiome matures sufficiently before age 1 year," says Jakob Stokholm, senior scientist at The Copenhagen Prospective Study on Asthma in Childhood and the study's first author. "Our study proposes the perspective of restoring a [caesarean section](#)-perturbed microbiome and thereby perhaps prevent [asthma](#) development in a child, who is otherwise at high risk."

More information: "Delivery mode and gut

microbial changes correlate with an increased risk of childhood asthma," *Science Translational Medicine* (2020). [stke.sciencemag.org/lookup/doi ... 26/scisignal.aax9929](https://stke.sciencemag.org/lookup/doi/10.1126/scisignal.aax9929)

Provided by Rutgers University

APA citation: The hidden reason children born by C-section are more likely to develop asthma (2020, November 11) retrieved 27 April 2021 from <https://medicalxpress.com/news/2020-11-hidden-children-born-c-section-asthma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.