


3-D printers may be toxic for humans

December 15 2020



Credit: CC0 Public Domain

Risk researchers are asking new questions about the health and safety implications and how to mitigate any potential health risks to users of 3-D printers and consumers of products manufactured with this emerging technology, especially children. Such printing is increasingly being used in homes, schools, libraries and other spaces where people

commonly spend a lot of time. The particles released during the printing process, which are small enough to infiltrate deep into the lungs, can affect indoor air quality and public health. The wide use of 3-D printers to manufacture face shields, respirators and other personal protective equipment for COVID-19 has created a new urgency on these questions.

Several studies that aim to characterize and quantify the release and composition, [particle size](#), and residence time in the indoor environment will be presented in the Exposure and Risk Assessment of 3-D Printing and Emerging Materials symposium on December 15 at the 2020 Society for Risk Analysis virtual Annual Meeting held December 13-17, 2020.

The base materials used in 3-D printers include thermoplastics, metals, nanomaterials, polymers and volatile and semi volatile organic chemicals. The [printing process](#) may take several hours, and during this time a range of chemical by-products and particulates may be released into indoor environments.

Given these unknowns, scientists have begun to conduct studies to understand these releases and their specific composition, particle size, and residence time in the indoor environment, producing data that can be incorporated into robust exposure and risk assessments.

A study conducted by Yong Qian, Ph.D., National Institute for Occupational Safety and Health (NIOSH), evaluated the potential toxicity of ABS emissions generated during 3-D printing by examining human lung cells and rats exposed via inhalation. The study, "Acrylonitrile butadiene styrene (ABS) printer emission induced in vitro and in vivo toxicity," revealed that the emitted particles cause moderate toxicity in human lung cells and minimal toxicity in rats.

The presentation, "Recent 3-D printing emissions research at Environmental Protection Agency (EPA)," reviews two recent studies

from the EPA; the first analyzed emissions from a 3-D printer filament extruder (a device used to create 3-D printer filaments) in a laboratory setting, and the second used a simulation model to predict the number of particles deposited at specific locations in the respiratory tract, and how that pattern changes for individuals of different ages, when using a 3-D printer.

"To date, the general public has little awareness of possible exposures to 3-D printer emissions," states Peter Byrley, Ph.D., EPA, lead author. "A potential societal benefit of this research is to increase public awareness of 3-D [printer](#) emissions, and of the possibly higher susceptibility of children."

The studies found that the filament extruder released amounts of small particles and vapors similar to those found in studies of 3-D printers, and the simulation model predicted higher deposition of particle mass per surface area in the pulmonary region for individuals ages nine and younger. Further testing of emission profiles with additional simulation studies to predict inhaled dose are needed.

While 3-D printing makes numerous products more readily available, and at cheaper costs, they do contribute to the mass of plastic products polluting the planet. Joana Marie Sipe, Duke University, has developed a machine that can measure how much a plastic product, like a water bottle, can break down through rubbing and sanding during use and in the environment. The plastic particles were then fed to fish to see the effects that the nanoparticles in the plastic had on their organs.

The study, "NanoPHEAT: Forecasting nanocomposite consumer product's release, exposure, and toxic effects of nanomaterials (MWCNT and Ag NPs)," reveals that when plastics breakdown, the nanomaterials that were incorporated and thought to be biologically unavailable become exposed to the environment. The researchers were

able to predict the percentage of nanoparticles that came out of the plastic when they were eaten by the fish, providing a Matrix Release Factor (MRF) which could be used to find out the quantity of plastic and nanoparticles that are released when someone chews a product or when it breaks down in the ocean.

"This research can help set regulations on how much nanomaterial fillers can be added to particular consumer products, based on their MRF value," states Sipe. "The data can help determine how much plastic and/or nano-filled products release contaminants into the environment or the human body."

As 3-D printing technologies become more widespread, regulators, manufacturers, and users may need to focus their attention on better managing potential risks.

Provided by Society for Risk Analysis

Citation: 3-D printers may be toxic for humans (2020, December 15) retrieved 10 January 2023 from <https://medicalxpress.com/news/2020-12-d-printers-toxic-humans.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--