


Prediabetes subtypes identified

4 January 2021



In people with prediabetes, there are six clearly distinguishable subtypes (clusters), which differ in development of the diseases, risk for diabetes, and development of complications. Credit: IDM

All prediabetes is not the same: in people in the preliminary stages of type 2 diabetes, there are six clearly distinguishable subtypes, which differ in the development of the disease, diabetes risk, and the development of secondary diseases. This is shown in a study by the Institute for Diabetes Research and Metabolic Diseases (IDM) of Helmholtz Zentrum München; at the University of Tübingen, Tübingen University Hospital; and the German Center for Diabetes Research (DZD). The results have now been published in *Nature Medicine*. The new classification can help in the future to prevent the manifestation of diabetes or the development of diabetes complications through targeted prevention.

Diabetes is a worldwide pandemic. Since 1980, the number of people with diabetes has quadrupled worldwide. In Germany alone, 7 million people suffer from it. And the trend is still rising. By 2040, the number of people with type 2 diabetes could increase to as many as 12 million. But type 2 diabetes does not develop from one day to the next. People often go through a longer preliminary stage of diabetes, in which blood glucose levels

are already elevated but people are not yet ill. "For people with prediabetes it has not been possible until now to predict whether they would develop diabetes and be at risk for serious complications such as kidney failure, or whether they would only have a harmless form with slightly higher blood glucose levels but without significant risk," said Professor Hans-Ulrich Häring, who initiated the study 25 years ago.

However, such a distinction is important for the targeted prevention of the metabolic disease and thus for counteracting the diabetes pandemic. Researchers from Tübingen have now achieved an important breakthrough. Using [cluster analysis](#) in people with prediabetes, they have discovered six distinct subtypes with different [diabetes risk](#). A differentiated classification of prediabetes and diabetes makes it possible to carry out individual and early prevention and therapy of diabetes and its secondary diseases in a way that is adapted to the development of the disease.

The research group led by Professor Häring and Professor Fritsche at the University Hospital in Tübingen has conducted detailed studies of the metabolism of people with prediabetes who are still considered healthy. The test persons (n=899) are from the Tübingen Family Study and the study of the Tübingen Lifestyle Program. They have repeatedly undergone intensive clinical, laboratory chemical, magnetic resonance imaging and genetic examinations in Tübingen over the past 25 years.

Based on key metabolic parameters such as blood glucose levels, liver fat, body fat distribution, blood lipid levels and genetic risk, the researchers were able to identify six subtypes of prediabetes. "As in manifest diabetes, there are also different disease types in the preliminary stage of diabetes, which differ in [blood glucose levels](#), insulin action and insulin secretion, body fat distribution, liver fat and genetic risk," said first author Professor Robert Wagner from the DZD Institute for Diabetes Research and Metabolic Diseases (IDM) of Helmholtz Zentrum München at the University of

Tübingen, summarizing the results of the study.

Three of these groups (clusters 1, 2 and 4) are characterized by a low diabetes risk. The study participants in clusters 1 and 2 were healthy. Slim people are the main members of cluster 2. They have a particularly low risk of developing complications. Cluster 4 consists of overweight people, whose metabolism is still relatively healthy. The three remaining subtypes (clusters 3, 5 and 6) are associated with an increased risk of diabetes and/or secondary diseases. People who belong to subtype 3 produce too little insulin and have a high risk of developing diabetes. People in cluster 5 have a pronounced fatty liver and a very high risk of diabetes because their bodies are resistant to the blood glucose lowering effect of insulin. In subtype 6, damage to the kidneys occurs even before diabetes is diagnosed. Here, mortality is also particularly high.

But can the classification into six prediabetes subtypes also be confirmed in other cohorts? To investigate this, the researchers extended the analysis to include almost 7000 subjects in the Whitehall II Cohort in London and there, too, identified the six prediabetes subtypes.

The researchers are already making further plans. "Next, in prospective studies, we will first seek to determine to what extent the new findings are applicable for the classification of individual persons into risk groups," said Professor Andreas Fritsche of Tübingen University Hospital. If this is the case, people with a high risk profile could be identified early on and receive specific treatment.

These results are based on research conducted by the Diabetes Research Department at Tübingen University Hospital over the past 25 years to characterize people with an increased risk of diabetes. The study was funded by the German Research Foundation, the Federal Ministry of Education and Research and the State of Baden-Württemberg.

"One of the DZD's goals is to develop precise prevention and therapy measures, i.e. the appropriate prevention or treatment for the right group of people at the right time. The combination

of in-depth clinical and molecular research with state-of-the-art bioinformatics has made this internationally important result possible. The identification of subtypes in the preliminary stages of type 2 diabetes is an important step towards precision medicine in the prevention of [diabetes](#) and its complications," says DZD Executive Director Prof. Martin Hrab? de Angelis.

More information: Robert Wagner et al. Pathophysiology-based subphenotyping of individuals at elevated risk for type 2 diabetes, *Nature Medicine* (2020). [DOI: 10.1101/2020.10.12.20210062](#)

Provided by Deutsches Zentrum fuer Diabetesforschung DZD

APA citation: Prediabetes subtypes identified (2021, January 4) retrieved 12 June 2022 from <https://medicalxpress.com/news/2021-01-prediabetes-subtypes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.