

Teen vaping doubles risk for subsequent tobacco use

7 January 2021



current (past 30 days) <u>cigarette smokers</u> (RR, 2.20) and SLT users (RR, 1.64).

"It is important to remain vigilant in seeking strategies and policies to prevent teens from starting habits that pose a threat to their health, particularly when they serve as a stepping stone to more harmful tobacco products," Keller-Hamilton said in a statement.

More information: Abstract/Full Text (subscription or payment may be required)

Copyright © 2020 HealthDay. All rights reserved.

(HealthDay)—Adolescent boys who use electronic cigarettes have an increased risk for subsequently initiating traditional tobacco products, according to a study published online Dec. 8 in *Addictive Behaviors*.

Brittney Keller-Hamilton, M.P.H., from the Comprehensive Cancer Center at The Ohio State University in Columbus, and colleagues used a causal inference framework to identify whether male adolescent e-cigarette users were at increased risk for initiating cigarette smoking and smokeless tobacco (SLT) use compared with similar boys who had never used e-cigarettes. The analysis included 1,220 boys (aged 11 to 16 years at enrollment) followed every six months for two years.

The researchers found that compared with nonusers, e-cigarette users were more than twice as likely to later initiate both cigarette smoking (relative risk [RR], 2.71) and SLT (RR, 2.42). E-cigarette users were also more likely to become



APA citation: Teen vaping doubles risk for subsequent tobacco use (2021, January 7) retrieved 25 July 2022 from https://medicalxpress.com/news/2021-01-teen-vaping-subsequent-tobacco.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.