


Approximately half of AD dementia cases are mild, one-fifth are severe

13 January 2021



Credit: CC0 Public Domain

What percent of patients with Alzheimer's Disease (AD) currently have severe dementia? Do more people have mild disease? Or are the majority suffering with moderate dementia? A new study using data from the Framingham Heart Study (FHS) sheds light on these trends. Boston University School of Medicine researchers have found that slightly more than half (50.4 percent) of cases are mild, just under one-third (30.3 percent) of cases are moderate and 19.3 percent are severe cases. Among all participants with mild cognitive impairment (MCI) and AD, the pooled percentage was 45.2 percent for the combined group of mild AD dementia and MCI that later progressed to AD.

"Early intervention in MCI or the mild stage of AD dementia has been the primary focus for AD research and [drug development](#) in recent years. We found that approximately 45 percent of all those who are cognitively impaired or diagnosed with AD-dementia had early AD. Our results serve to inform the design of future research studies such as clinical and [observational studies](#) and provide optimal resource allocation for policy-

making," explained corresponding author Rhoda Au, Ph.D., professor of anatomy and neurobiology at Boston University School of Medicine.

To characterize the distribution of severity of AD dementia and MCI among prevalent cases in the population, FHS participants (aged 50-94) with prevalent MCI or AD dementia clinical syndrome were selected from three time-windows: 2004-2005, 2006-2007 and 2008-2009. Estimates of the severity distribution were achieved by pooling results across time-windows. Diagnosis and severity were assessed by consensus dementia review. MCI-progressive was determined if the participant had documented progression to AD dementia clinical syndrome using longitudinal data.

According to the researchers the finding that half of the people living with AD have mild disease underscores the need for research and interventions to slow decline or prevent progression of this burdensome disease. "It is crucial to determine [risk factors](#) or develop therapies that could alter the disease trajectory to improve individuals' quality of life and alleviate the socio-economic burden," adds Au.

The researchers believe that most people who have AD are still at a stage when there is still some preserved quality of life. "This means any drug treatment that is effective might help prevent their AD from getting worst."

More information: Jing Yuan et al, Severity Distribution of Alzheimer's Disease Dementia and Mild Cognitive Impairment in the Framingham Heart Study, *Journal of Alzheimer's Disease* (2020). [DOI: 10.3233/JAD-200786](https://doi.org/10.3233/JAD-200786)

Provided by Boston University School of Medicine

APA citation: Approximately half of AD dementia cases are mild, one-fifth are severe (2021, January 13) retrieved 11 September 2022 from <https://medicalxpress.com/news/2021-01-approximately-ad-dementia-cases-mild.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.