

COVID lockdown loneliness linked to more depressive symptoms in older adults

January 22 2021



Credit: CC0 Public Domain

Loneliness in adults aged 50 and over during the COVID-19 lockdown was linked to worsening depressive and other mental health symptoms, according to a large-scale online study.



Loneliness emerged as a key factor linked to worsening symptoms of depression and anxiety in a study of more than 3,000 people aged 50 or over led by the University of Exeter and King's College London, and funded by The National Institute for Health Research (NIHR) Maudsley Biomedical Research Centre (BRC).

Researchers had access to data going back to 2015 for participants of the PROTECT <u>online study</u>. They also found that a decrease in <u>physical</u> <u>activity</u> since the start of the pandemic was associated with worsening symptoms of depression and anxiety during the pandemic. Other factors included being female and being retired.

Dr. Byron Creese, of the University of Exeter Medical School, who led the study, said: "Even before the pandemic, loneliness and physical activity levels were a huge issue in society, particularly among older people. Our study enabled us to compare mental health symptoms before and after COVID-19 in a large group of people aged 50 and over. We found that during lockdown, loneliness and decreased physical activity were associated with more symptoms of poor mental health, especially depression. It's now crucial that we build on this data to find new ways to mitigate risk of worsening mental health during the pandemic."

The study found that before the pandemic, lonely people would report an average of two symptoms of depression for at least several days over the previous last two weeks. During lockdown, lonely people reported either an increase in frequency of depressive symptoms, to more than half the days in the two week period, or a new symptom for at least several days in that timeframe. In people who were not lonely, levels of depressive symptoms were unaffected.

PROTECT began in 2011, and has 25,000 participants signed up. Designed to understand the factors involved in healthy ageing, the innovative study combines detailed lifestyle questionnaires with



cognitive tests that assess aspects of brain function including memory, judgment and reasoning over time. In May, researchers included a new questionnaire designed to assess the impact of COVID-19 on health and wellbeing. Running from May 13 to June 8, the questionnaire was completed by 3,300 people, of which 1,900 were long-standing PROTECT participants. The study is continuing to run so that longer term outcomes can be assessed.

Zunera Khan, Research Portfolio Lead at Institute of Psychiatry, Psychology & Neuroscience said "We've found links between loneliness and a drop in physical exercise and worsening mental health symptoms. It should be within our power to find ways of keeping people socially engaged and active. Our online PROTECT platform ultimately aims to find new ways to engage people in their homes, however, technology can only be part of the picture. We need to ensure we can find new ways to help people stay active and social, whether they are online or not."

Professor Clive Ballard, Executive Dean and Pro-Vice Chancellor of the University of Exeter Medical School, who leads PROTECT, said: "We are only just beginning to learn the impact that COVID-19 is having on the health and wellbeing of older people. For example, the effect of any economic impact may not yet have emerged. Our largescale study will span a number of years, and will help us understand some of the longer-term effects of COVID-19 on mental health and wellbeing, and ultimately, on whether this has any knock-on effect on aspects of ageing, such as brain function and memory."

The study plans to conduct further analysis on groups at particularly high risk, such as people with cognitive impairment and those with caring roles

More information: Byron Creese et al. Loneliness, physical activity and mental health during Covid-19: a longitudinal analysis of depression



and anxiety in adults over 50 between 2015 and 2020, *International Psychogeriatrics* (2020). DOI: 10.1017/S1041610220004135

Provided by University of Exeter

Citation: COVID lockdown loneliness linked to more depressive symptoms in older adults (2021, January 22) retrieved 9 March 2023 from https://medicalxpress.com/news/2021-01-covid-lockdown-loneliness-linked-depressive.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.